Plant Disease Reporter

Agricultural Research Center

Agricultural Research Service

UNITED STATES DEPARTMENT OF AGRICULTURE

VOLUME 58 NUMBER 13

PLDRA4 58(13) 1145-1186

1974 Index
(Issued November 1975)

The Plant Disease Reporter is published monthly as a service to American agriculture. The articles emphasize new developments in plant pathology, including disease incidence and severity, serious outbreaks and epidemics, investigative techniques, and control chemicals. The rapid issuance of such information permits plant pathologists to recommend the judicious use of pesticides or other appropriate control measures against impending disease epidemics. Agricultural Research Center, in publishing this material, does not assume responsibility for the subject matter.

For sale by the Superintendent of Documents, U.S. Government Printing Office,
Washington, D.C. 20402 - Subscription Price: $22.40 per year;
$5.60 additional for foreign mailing. Single copies vary in price

Issued by

APPLIED PLANT PATHOLOGY LABORATORY
Plant Protection Institute

Beltsville Agricultural Research Center
Volume 58

Beltsville, Maryland
Number 13

AUTHOR INDEX

ABAWI, G. S., (774)
ABDALLA, NABIL, 562
ABDEL-RAHMAN, T. B., 517
ABDOU, Y. A., 176
ABNEY, T. S., (789)
ABOUL-EID, H. Z., 1130
ACCIAVATTI, ROBERT E., 418
ACOSTA, J. M., (122)
ADAMS, P. B., 700
AGRIOS, GEORGE N., (536)
ALBRECHTSEN, R. S., (743)
ALDWINCKLE, HERB S., 625, 696, 819
ALE-AGHA, N., 915, (983)
ALFARO, A., 549
ALLEN, D. J., 896
ALLEN, R. M., (945)
ALLEN, T. C., (269)
ALLEN, T. D., (793)
AMES, RONALD, (162)
AMIN, K. S., 358
AMMERMAN, BROOK A., (718)
AMOSU, J. O., 361
ANDERSON, F. N., (379), 782
ANDERSON, ROBERT L., 92
ARMSTRONG, G. M., 479
ARMSTRONG, JOANNE K., (479)
ARNETT, JAMES D., Jr., 754
ARTMAN, J. D., 409
ATHOW, K. L., (130), 789
ATIF, A. H., (1085)
AUGER, JAIME G., 599
AWODERU, V. A., (544)
AYERS, J. E., (934)

bALD, J. G., 126
BALSBAUGH, EDWARD U., Jr., (906)
BANIECKI, J. F., (291), 421
BARKER, K. R., (822), 991
BARKSDALE, T. H., 406
BARNES, G. L., 687
BARNETT, R. D., (3), (878)
BARRATT, D. H. P., (777)
BATCHELOR, D. L., 830
BATRA, L. R., 719
BAXTER, L. W., Jr., 139, 300, 844
BEGA, ROBERT V., (102), 1069
BeMILLER, J. N., (1113)
BERGDAHL, DALE, 616
BERGER, R. D., 922
BERGQUIST, R. R., 495
BERTUS, A. L., 437
BEUTE, M. K., (348)
BEUTEL, J. A., (99)
BILGRAMI, R. S., (1134)
BILLETER, B. A., (1065)
BIRD, G. W., 541
BISSONNETTE, HOWARD L., (693)
BISTLINE, F. W., (895)
BLACK, W. M., 511
BLACKMON, CYRIL W., 641
BLAINE, WILLIAM M., (448)
BLAIR, R. L., (198)
BLAKE, C. T., (822)
BLANCO, M. H., 934
BLUE, R. L., (757)
BOESEWINKEL, H. J., 525, 705
BOLKAN, H. A., 646
BOOTH, R. H., (703)
BOSE, ELAINE, (663)
BOSWELL, F. C., (541)
BOWEN, H. H., (780)
BRATHWAITE, CHELSTON W. D., 259, 1048
BRAUN, A. L., (633)
BRENER, W. D., 653
BROOKS, JAMES L., 291
BROOKS, O. L., (541)
BROWDER, L. E., (469)
BROWN, A. R., (21)
BROWN, G. ELDON, (933)
BRUEHL, G. W., 554
BRZOSTOWSKI, H. W., 92
BUFORD, W. R., (1143)
BURGIS, D. S., (579)
BURKE, D. W., (500)
BURNETT, H. C., 355
BUTTERFIELD, J. E., (367), (558)
BYTHER, R. S., 401

CAFLEY, J. D., (677)
CALAVAN, E. C., (82), (757), (850)
CALUB, A. G., 956
CANCINO, LAUTARO, 568
CANERDAY, T. D., (541)
CARLYSLE, THELMA C., (1051)
CARROLL, R. B., 443
CARTER, M. V., 442
CEPONIS, M. J., 367, 558
CHALFANT, R. B., (239)
CHALKLEY, JAMES H., (195)

CHAMBERLAIN, D. W., 50
CHANDLER, W. A., 208, 388
CHARUDATTAN, R., (277)
CHHEDA, H. D., (616)
CHILDS, J. F. L., 1051
CHORIN, MATHILDA (45)
CHOU, L. G., 221
CHRISTIE, S. R., 658
CLIFFORD, D. R., (949)
COBB, F. W., Jr., (478)
COHEN, MORTIMER, 801
COLE, H., Jr., (41), (136)
COMPTON, W. A., (956)
COMSTOCK, J. C., 104
CONNOR, JAMES F., (718)
COOK, G. E., (190)
COOK, R. J., (62)
COPPOLINO, JOHN B., (660)
CORBETT, M. K., (17)
CORDELL, CHARLES E., 188
CORLEY, W. L., (899)
COTHER, E. J., 1126
COVEY, RONALD P., Jr., (701)
COYNE, DERMOT P., 278, 379, (782)
CRAWFORD, JOHNNY L., (369), 373
CRILL, PAT, 579
CROSBY, F. L., (666)
CUNFER, B., 679

DABUSH, S., (1142)
DAINES, ROBERT H., (17), 254
DALMACIO, S., (934)
DANCE, B. W., 677
DANGE, S. R. S., 285
DAS, C. R., (358)
DAUBENY, HUGH A., 391, 1024
DAVIS, DONALD D., 14, 660
DAVIS, R. E., 1109
DAVIS, S. H., Jr., (387)
DAWSON, DAVID H., (721)
DEAN, C. E., (658)
DEAN, JACK L., 439
DEBROT, E. A., 1032
DEMSKI, JAMES W., 195, 723
DEWEY, W. G., 743
DHINGRA, O. D., 1034
DICKSON, D. W., (374)
DICKSON, M. H., 774
DILBECK, J. D., (374)
DINUS, RONALD J., 584
DOBSON, J. W., (247)
DOWLER, W. M., (76)
DRIVER, CHAS. H., (492)
DRUMMOND, D. B., (1105)
DUBOS, BERNADETTE, (71), 147, (915)
DuCHARME, E. P., (793)
DUNKLE, LARRY D., (267)
DuVALL, T. W., (521)
DWINELL, L. D., 198, 497

EBRON, T. T., Jr., (544)
EDMONDS, ROBERT L., 1077
EDMUNDS, L. K., (267)
EDWARD, J. C., (345)
EDWARDS, GEORGE J., 793
ELGINDI, D. M., (517)
ELLETT, C. W., (978)
ELLIS, M. A., 173, 332, 534, 760
ELMSTROM, G. W., (114)
ENGELHARD, ARTHUR W., 669
ENTEMANN, F. M., (62)
EPPS, W. M., (844)
EPSTEIN, A. H., (104)
ESCAFFI, O., (599)
ETHREDGE, W. J., (666)
EVANS, A. K., (110)
EVERSMEYER, M. G., 459, 469
EVERSON, E. H., (522)

FAGAN, SUSAN G., (139), (300)
FAJEMISIN, J. M., 313
FARLEY, J. D., 320
FASSULIOTIS, GEORGE, 1000
FATOKUN, D. O., (616)
FELDMAN, A. W., 35
FENSTERMACHER, JAMES M., (573), (848)
FISCHER, H. U., (141), 143, 252, (307), (309), 1117, 1121, 1123
FLAVELL, R. B., (777)
FORDE, H. I., (630)
FORER, L. B., 987
FRANCKOWIAK, J. D., (361)
FRAZIER, N. W., 28, 203
FREEMAN, JACK A., (391)
FREEMAN, T. E., 277
FRENCH, D. W., (616)
FREYTAG, R. E., (825)
FUDL-ALLAH, A. E.-S. A., 82
FUTRAL, J. G., (541)

GARDNER, C. O., (956)
GENTER, C. F., (456)
GENTNER, W. A., (165)
GHANEKAR, A. M., 354
GHORAB, A. I., (1130)
GILL, D. L., 710, 1012, 1015
GILMER, R. M., (616)
GIVEN, J. BRUCE, (291)
GOHEEN, A. C., (549)
GONSALVES, D., (895)
GOODING, G. V., Jr., 521
GOODMAN, R. N., (620)
GOUGH, F. J., 80
GRACE, T. D. C., (92)
GRAND, L. F., 318, (439)
GRANETT, A. L., (155), 211
GRAU, CRAIG, 693
GRAY, L. E., (50)
GREEN, DOV, (683)
GREGORY, GAROLD F., 65
GROSCLAUDE, C., 71, (915), (983)

GUMPF, D. J., (1002)
GUPTA, PRAMILA, 345
GUZMAN, V. L., (1087)

HAESELER, C. W., 486
HAGEDORN, D. J., 226, 229, (322), 330, (941)
HALLIWELL, ROBERT S., 23
HAMMOND, D. F., 17
HAMPSON, M. C., 971
HANKINS, B. J., (432)
HANKS, R. W., (35)
HARDER, H. H., (26)
HARE, ROBERT C., (584)
HARMON, S. A., (239), (746), (749)
HARRISON, R. L., (456)
HARVEY, I. C., 57
HEALD, C. M., (337), 985
HELBURG, L. B., (571)
HERR, L. J., 157
HERRON, J. W., (90)
HILL, J. P., (934)
HINZ, P., (860)
HISLOP, E. C., 949
HOCH, H. C., 941
HOCKETT, E. A., (679)
HODGES, CLINTON F., 448
HODGES, C. S., 282
HODGES, LARRY R., (88)
HOES, J. A., (311)
HOITINK, H. A. J., 650
HOOKER, A. L., (219), (313), 399, 909, 975
HOPKINS, D. L., 114
HORSFALL, J. G., 161
HORST, R. K., 814
HORTON, B. D., 553
HOWELL, W. E., 766
HUBBELING, N., (320)
HUBER, D. M., 432
HUGHES, I. W., (974)
HYRE, R. A., 297

ILYAS, M. B., (332), (760)
IMADA, JUN, (603)
INSERRA, R. N., 1075
ISHII, M., (495)

JABERG, CAROL, (320)
JACKSON, NOEL, 573, (848)
JAIN, K. L., (285)
JATALA, PARVIZ, 591
JENKINS, LEONARD E., (890)
JENSEN, HAROLD J., (128), 269, (591), 594
JODON, MILDRED H., (814)
JOHN, V. T., (856)
JOHNSON, ALBERT S., (1037)
JOHNSON, A. W., 239, 746, 749
JOHNSON, H. G., (363)
JOHNSON, P. W., 730
JONES, A. L., (243)
JONES, JOHN PAUL, (579), 636
JONES, THOMAS W., (65)

KANG, M. S., 1113
KARAYIANNIS, IRENE, (490)
KASIMATIS, A. N., (869)
KEENAN, JOHN G., 890
KEENEY, D. R., (304)
KESSLER, KENNETH J., Jr., 1042, 1044
KHADR, A. S., (176)
KIM, SOON KWON, 219
KINLOCH, R. A., (169)
KIRKHAM, M. B., 304
KISSLER, J. J., (869)
KISTLER, B. R., (287)
KLISIEWICZ, J. M., 926
KLOS, E. J., (180), (181)
KMETZ, KONRAD, 978
KNAUSS, J. F., (445), 1100
KNUTSON, DONALD M., 235
KO, W. H., (86)
KOEPSELL, P. A., (269)
KOLLAS, D. A., (40)
KOMMEDAHL, THOR, 363
KONTAXIS, D. G., 904
KRAFT, J. M., 62, 500
KRATKY, B. A., 86
KRESS, L., (14)
KUCHAREK, T. A., 763, 878
KUHLMAN, E. G., (282)
KUHN, C. W., (107)
KUIPER, J., 845

LACASSE, N. L., (41), (136)
LACY, M. L., 232
LAEMMLEN, FRANKLIN, 102
LANA, ALLAN O., 536, 616, 1040
LANDIS, T. D., 110
LANE, CARL L., 67
LAPP, N. A., (318)
LARACH, WILFREDO, (568)
LARSEN, P. O., (334)
LATORRE, BERNARDO, (568)
LAUNDON, G. F., (423)
LAUT, JOHN G., 571
LAVIOLETTE, F. A., (130), (789)
LEAR, B., (562)
LELEJI, O. I., (1080)
LENTZ, P. L., 165
LEONARD, C. D., 918
LEONARD, K. J., 529, 532, 612
LEONE, IDA A., 683
LIGHTFIELD, JOHN W., (810)
LIM, S. M., (219), 811
LINZON, S. N., (1105)
LIPTON, WERNER J., (118)
LITTRELL, R. H., (1017)
LOCKHART, B. E. L., 141, (143), (252), 307, 309, (1117), (1121), (1123)
LORBEER, J. W., (837)

LOUIE, RAYMOND, 733
LOWNSBERY, B. F., 630, o33
LOZANO, J. C., (703)
LUCAS, L. T., 822
LUEPSCHEN, N. S., 26, (243)
LUKE, H. H., 3
LUMSDEN, R. D., (700)

MABRY, JAMES E., 810
MACHADO, C. C., (173), (411)
MACHTMES, ROLLIN, (6), (554)
MAGASI, LASZLO P., 892
MAGIE, R. O., 256
MAHMOOD, M., (32), (265)
MAI, W. F., (1007)
MAJORANA, G., 691
MAKKOUK, K. M., 1002
MALEK, R. B., 997
MALOY, OTIS C., 6, 10, 902
MAMLUK, OMAR F., 996
MANJI, B. T., (663)
MANLEY, JANET M., (892)
MANNING, W. J., 483
MARAMOROSCH, K., (96), (797)
MARLATT, R. B., 445
MARTIN, G. C., (630)
MARTIN, M. W., (965)
MARTINSON, C. A., (104)
MATHRE, D. E., (679)
MATHUR, S. B., (507)
MATUSZEWSKI, MARK, (188)
McCLANAHAN, R. J., (730)
McDANIEL, M. E., (80)
McDONALD, R. E., 1143
McFEELEY, JAMES C., 773
McILVEEN, W. D., (1105)
McINTOSH, A. H., (797)
McKENZIE, D. L., (342)
McVEY, D. V., (608)
MEISTER, H., (904)
MENGES, R. M., (985)
MERRILL, W., 287
MEYER, W. A., (514)
MILBRATH, GENE M., 420
MILLER, D. L., 275
MILLER, DOUGLAS R., 94
MILLER, P. M., 708
MILNE, K. S., (646)
MINK, G. I., (766)
MIRCETICH, S. M., (99)
MOHAMMAD, ABU, 32, 265
MOLLENHAUER, H. H., (780)
MOLLER, W. J., 99, (442), 869
MOODY, E. H., (630), (633)
MOORE, HOWARD D., (890)
MOREY, D. D. (21)
MORGAN, PAUL D., 492
MORRALL, R. A. A., 342
MOTSINGER, RALPH E., 369
MOYER, JAMES W., 41, 136
MUEHLBAUER, F. J., (62)
MUELLER, W. C., 848
MUKHOPADHYAY, A. N., 952
MULLER, IRMGARD, (472)
MUMFORD, PAULINE, (1113)
MURPHY, JUDITH A., (1113)
MUSEN, HAROLD L., (641)

NAGAICH, B. B., 968
NAUER, E. M., (757)
NEELY, DAN, 261, (511)
NEERGAARD, PAUL, (507)
NELSON, L. R., 21
NELSON, R. R., (934)
NGUNDO, B. W., 1020
NICHOLLS, THOMAS H., 1094
NICHOLS, LESTER P., 814
NICHOLSON, R. L., (430)
NOME, F. S., (599)
NORGREN, R. L., (653)
NORTON, D. C., (860)
NWIGWE, CHINENYE, 414, 416

O'BANNON, J. H., 561, (1075)
ODVODY, GARY N., 267
OFONG, A. U., 463
OGAWA, J. M., (663)
OGBUJI, R. O., 128, 594
OLIVEROS, B., 703
O'NEIL, R. T., (814)
ORDOSGOITTI, A., (1032)
ORMROD, D. J., (840)
ORR, CALVIN C., 120
OSHIMA, NAGAYOSHI, (890)
OTEIFA, B. A., (517)
OTTA, J. D., 476, 1061
OU, S. H., 544

PAGUIO, O. R., 107
PALM, H. L., (760)
PALTI, J., 45
PAPPELIS, A. J., (1113)
PARISH, C. L., 701
PARKER, K. G., 243, (784), 1007
PARMETER, JOHN R., Jr., (478)
PARTRIDGE, A. D., 75, (275)
PEARSON, R. G., 1105
PENROSE, L. J., 832
PEPIN, H. S., (391), 840, (1024)
PERIES, O. S., 293, 295
PERRY, C. E., (541)
PERRY, R. L., 780
PETERSEN, D. H., (486)
PETERSON, C. J., Jr., (554)
PETERSON, J. L., (387)
PETERSON, P. A., 777
PFLEGER, F. L., 1028
PHELPS, WILLIAM R., 596, 1137
PHILLEY, GEORGE, (23)
PHILLIPS, DOUGLAS J., 118
PHILLIPS, D. V., (723)
PIECZARKA, D. J., 837

PIIRTO, DOUGLAS D., 478
PINCKARD, J. A., (201)
PIÑERO, A., (122)
PINKAS, Y., (45)
PORTO, M. D. M., 322
POWELL, C. C., (911)
POWERS, H. R. Jr., (497)
PRASAD, S. S., 1134
PRASADA RAO, R. D. V. J., 856
PRASARTSEE, C., (173), (411)
PREY, ALLEN J., (1094)
PRINCE, V. E., (553)
PROVVIDENTI, R., 155, (211), 735
PURCIFULL, D. E., (658), (830)

RAHEJA, A. K., 1080
RAND, R. E., (229)
RAO, S. V. R. K., (952)
RASKI, D. J., (562)
RATHORE, R. S., (285)
RAVENSCROFT, A. V., (522)
REEDER, WILLIAM F., (59)
REINKE, ALEX W., (718)
REXRODE, CHARLES O., 382
RHOADS, ANN F., 102
RICARD, J. L., (71), (147), (915), 983
RICH, SAUL, 162, (428), (708)
RIDINGS, W. H., (355)
RIEDEL, R. M., 911
RÍORDÁIN, F. Ó., 12
RITCHIE, D. F., 181
ROANE, C. W., 455, 456
ROANE, MARTHA K., (455)
ROBERTS, EVAN P., (773)
ROBERTSON, L. W., (165)
ROBINSON, R. W., (735)
ROCHOW, W. F., 472
ROELFS, A. P., 605, 608, 806
ROHRBACH, K. G., (26)
ROISTACHER, C. N., 757, 850
ROTHMAN, PAUL G., 467, (605)
ROWE, R. C., 348
RUPPEL, E. G., 434
RYDER, EDWARD J., 1037

SAFFORD, JOANN, (334)
SAHARAN, G. S., 1030
SANBORN, R. R., (99)
SANDS, D. C., 40
SANTAMOUR, FRANK S., Jr., 727
SARAVANAPAVAN, T. V., (928)
SASSER, M., (443)
SCARBOROUGH, BETSY, (814)
SCHABER, BURTON D., 906
SCHENCK, N. C., 169
SCHIPPER, ARTHUR L., Jr., 721
SCHMIDT, R. A., 451
SCHMITT, C. G., 825
SCHMITT, D. P., 860
SCHMITT, R. V., (562)
SCHMITTHENNER, A. F., (221), (650), (978)
SCHOENEWEISS, DONALD F., 937
SCHOMAKER, M. E., (571)
SCHUSTER, M. L., (278), (956)
SCHWARTZ, H. F., (782)
SCHWENK, F. W., 352, (354)
SCIUMBATO, G. L., 201
SCOTT, D. H., (430)
SCOTT, H. A., 191
SCOTT, P. R., (434)
SETLIFF, E. C., (653)
SHAFFER, W. H., Jr., 620
SHANMUGANATHAN, N., 928
SHARMA, B. D., (358)
SHARMA, N. D., 640
SHARMA, R. K., (904)
SHILLINGFORD, C. A., 214
SHIVELY, OWEN D., 566
SHRINER, D. S., 439
SIEVERT, R. C., 1073
SIM, T., (352)
SIMS, J. J., (82)
SINCLAIR, J. B., (173), (332), (411), (534), (760), (1034)
SINCLAIR, W. A., (387), 784, (971)
SINGH, N. D., 122
SINHA, R. C., (968)
SIRADHANA, BABU SINGH, (285)
SISSON, M. A., (26)
SKELLY, JOHN M., (150), 396, 738
SKIDMORE, E. L., (459)
SKILLING, DARROLL D., 853, (1094), 1097
SKOTLAND, C. B., (10)
SKOVMAND, B., (1085)
SKOWRONSKI, B. S., 797
SLEESMAN, JOHN P., 334
SMITH, D. H., 666
SMITH, M. L., (1091)
SMOOKLER, M., 1142
SMOOT, J. J., 933
SOBERS, E. K., (1015), 1017
SONODA, R. M., 490
SOWELL, GROVER, Jr., 899
SPALDING, DONALD H., 59
SPIERS, A. G., (423)
STARLING, T. M., (455)
STEADMAN, J. R., 190, (379), (782)
STEELE, ARNOLD E., 88
STEINER, G. W., (401)
STRIDER, D. L., 875
STUDT, H. G., (133)
SUGGS, EDNA G., (612)
SUZUKI, TERRY, (126)
SZKOLNIK, MICHAEL, 326, 587

TAINTER, F. H., 622
TANAKA, HIROYASU, 603
TATE, C. J., (700)
TATE, K. G., 663
TAYLOR, D. P., (1020)
TAYLOR, JACK, 247

TENNE, F. D., 411, (760)
THAMES, W. H., 337
THANKAMMA, L., 767
THOMAS, P. E., 965
THOMPSON, SAMUEL S., (369)
THOMSON, S. V., 945
TIMMER, L. W., 504, 882
TINER, JAMES, (622)
TOSIC, M., (420)
TOWERS, BARRY, 718
TRUJILLO, G. E., 122
TUCKER, D. P. H., (355), 895
TUCKER, MELVIN, (622)
TURGEON, A. J., 514
TURNER, C. E., (165)
TYLER, L. J., (784)

UNDERHILL, E. M., (451)

VAKILI, NADER G., 96
Van ALFEN, NEAL K., 924
van KRAAYENOORD, C. W. S., 423
Van SCYOC, C. E., (191)
Van SCYOC, J. V., (191)
Van SICKLE, G. A., 872
VARDARO, P. M., (483)
VASILOFF, G. N., 1091

WADDELL, CLINTON D., (1097)
WADE, E. K., (330)
WAGNER, V. R., (987)
WALKER, J. H., (763), (878)
WALLIN, EUNICE K., (492)
WALTERS, JAMES W., 1066
WALTON, GERALD S., 428, (924)
WANG, ERNIE, L. H., 974
WARREN, H. L., 430
WASHBURN, BROOKS, (622)
WAYLAND, J. R., (985)
WEAVER, D. J., 76
WEBER, P. V. V., 387
WEHUNT, E. J., (76), (553)
WEINGARTNER, D. P., 180, 374
WEIR, L. C., (75)
WEISS, MELVYN J., (418)
WELCH, D. S., (784)
WELLS, J. C., (348)
WELTZIEN, H. C., 133
WENHAM, H. T., (646)
WHITESIDE, J. O., 713
WICKS, T. 886
WIESE, M. V., 522
WILCOX, J. R., 130
WILCOXSON, ROY D., 1085
WILFRET, G. J., (256)
WILL, J. B., 150
WILLIAMS, A. S., 90
WILLIAMS, ERVIN, Jr., 272
WILLIAMS, H. E., (99)
WILLIS, WILLIAM G., 566
WILLSON, HOWARD, (10)
WINDELS, CAROL E., (363)
WINFIELD, A. L., 913
WINTER, WALTER E., 507
WITCHER, WESLEY, (67), (754)
WOLF, E. A., (922)
WOLTZ, S. S., 184
WOOD, FRANCIS A., (738)
WRIGHT, W. R., 1065
WUEST, P. J., (987)
WYNNE, J. C., (348)

YARWOOD, C. E., 54, 490, (733)
YOUNG, H. C., Jr., (272)
YOUNG, J. B., (521)
YOUNG, R. J., (291), (421)
YOUNG, T. R., (830)

ZAHNER, S. A., 793
ZEHR, E. I., (844)
ZIMMER, D. E., 311
ZITTER, T. A., 1087
ZUKEL, J. W., (162)

SUBJECT INDEX

Abelmoschus esculentus (see Okra)
Abies concolor (white fir): a tip dieback caused by Sydowia polyspora found for first time on white fir and for the first time in California, 94
--- fraseri (Fraser fir): root rot caused by Phytophthora cinnamomi, has become an increasingly important disease of Fraser fir Christmas trees in North Carolina, 318
--- grandis (grand fir): root rot indicators, primarily for Armillaria mellea, are bark beetle galleries or discolored foliage, 275
Acaulospora laevis: 169
Acrocylindrium oryzae: 358
Adjuvants: Nu-Film 17 and Nu-Film P had little effect on efficacy of Dyrene or chlorothalonil against Septoria apiicola on celery, 232
Aechmea fasciata: Helminthosporium rostratum is cause of new foliar disease of this ornamental, 445
Africa: 896
Air pollution (see also Ozone): relative susceptibility of 10 bean varieties to ozone, 14; ozone injury of Poa annua reduced with benomyl and thiophanate, 41; naturally occurring oxidant injury on azalea plants was suppressed with

benzimidazole or oxathiin compounds, 136; high nitrogen fertilizer in fall reduced air pollution symptoms in eastern white pine Christmas trees in Virginia, 150; potato plants grown in a growth chamber without filtered air are subject to air pollution damage, 304; relationship between age and ozone sensitivity of current needles of ponderosa pine, 660; a photocopy technique to evaluate fluoride injury on gladiolus in Ontario, 1091; injury on tomatoes in southwestern Ontario; PAN is the most likely cause, 1105

Alfalfa: soil pH had little influence on susceptibility to Meloidogyne hapla of resistant or susceptible varieties, 594

Almond (Prunus amygdalus): newly recognized leaf scorch disease of almond found in California, 99

Alternaria spp.: 130; 180; 332

--- macrospora: 201

--- tenuis: 558

Ambrosia beetle (Xylosandrus germanus): 1044

Anacardium occidentale (cashew): Phytophthora nicotianae var. nicotianae was isolated from cashew trees with shoot rot and leaf fall symptoms, 767

Antagonism: of Gliocladium roseum to Eutypa armeniacae, the pathogen that causes apricot canker, 983

Anthracnose: of pea, first epiphytotic in Wisconsin since 1942, 226; anthracnose and stalk rot of corn caused by Colletotrichum graminicola, 532

Antibiotics (see also Chemicals, Fungicides and Nematicides): streptomycin sulfate was phytotoxic to soybean seedlings, thus is not useful to control bacterial diseases of soybean as a soil drench, 534; twice as much streptomycin in concentrated spray (low volume) was absorbed by apple foliage than with dilute spray, but phytotoxicity occurred and apple fruits were reduced in size, 620; tetracycline drenches applied to young tree-declined citrus trees resulted in improvement in three of five trees treated, 895; Lowden formulation containing nystatin for control of Dutch elm disease was found to be effective if inoculation of elms and treatment with nystatin occurred the same day, 924

Antirrhinum (snapdragon): powdery mildew (Oidium sp.) controlled with BAS 67054 F, HOE 2873, thiophanate-methyl and triarimol fungicides, 12

Aphanomyces cochlioides: 157

Aphelenchoides fragariae: 911

Aphelenchus spp.: 1048

Aphis craccivora: 1080

Apple: minimum rates of pesticides on apples is dependent upon the amount of materials needed per acre for pest control regardless of the amount of water used, 247; low volume (concentrated) spray of streptomycin resulted in greater uptake of the antibiotic than dilute spray (high volume) but caused some phytotoxicity, 620; control of aerial galls with 2,4-xylenol and m-cresol, 701; eradication of powdery mildew from apple buds with methyl alkanoates and aliphatic alcohols, 949; damage from Pratylenchus penetrans on different root stocks, 1007

---: cedar-apple rust: an emulsifiable concentrate formulation of triforine controlled cedar-apple rust in greenhouse tests on 'Rome Beauty' with single dilute spray 1, 2, or 3 days after inoculation, 587; susceptibility of 41 apple cultivars to Gymnosporangium juniperi-virginianae and G. clavipes, 696

---: fire blight: a laboratory method described for testing pathogenicity of suspected Erwinia amylovora isolates, 181; inoculations were made with E. amylovora to young apple rootstock selections to record infection progress downward in the inoculated twig, 243; field susceptibility of 46 apple cultivars to fire blight, caused by E. amylovora, 819

---: scab: use of a combined temperature-leaf wetness recorder for improved apple scab control in Lebanon, 133; tolerance of apple scab fungus, Venturia inaequalis, to benomyl and thiophanate-methyl, 886

Apricot: Eutypa armeniacae was associated with pruning wound cankers on apricot trees in Spain, 442

Arachis hypogaea (see Peanut)

Arceuthobium spp.: 235

--- americanum: 235; 1066

--- microcarpum: 418

Argentina: 507

Arizona: 418

Arkansas: 622

Armillaria mellea: 102; 275; 719

Artichoke: 691; an artichoke virus found in Morocco is considered to be a variant of artichoke mottled crinkle virus, 1117; artichoke latent virus was found to be widespread in commercially-grown artichokes in Morocco, 1123

Ascochyta gossypii: 373

--- graminea: 455

--- rabiei: first record of Ascochyta rabiei,

a destructive pathogen of chickpea, in North America, in Saskatoon, Canada, 342
--- sorghi: 455
Ash (see Fraxinus)
Asparagus: Phytophthora megasperma var. sojae is recorded for the first time in New Zealand causing a new disease of asparagus, 525
Aspen (see Populus)
Aspergillus spp.: 332
--- flavus: 1134
--- melleus: 332
--- nidulans: 1134
--- niger: 1134
--- quadrilineatus: 1134
--- variecolor: 1134
Assay: of soil for Fusarium soilborne inoculum potential for gladiolus corm rot, 184; biochemical assay with Helminthosporium maydis race T pathotoxin to detect "Texas" cytoplasm in corn varieties, 777; seed assay for detection of Verticillium in safflower seed, 926
Aster (Michaelmas daisy): new wilt of aster in New Zealand is caused by Phialophora fastigiata, 57
Aureolaria grandiflora var. serrata ("false foxglove"): was found parasitizing eastern redcedar in Texas, 773
Australia: 92; 437; 845; 886; 1126
Azalea (Rhododendron kurume): oxidant injury on rooted cuttings suppressed by benomyl drenches and foliar sprays of thiophanate, 136

Baby's breath (Gypsophila paniculata): serious new crown rot and wilt found in Florida, caused by Phytophthora parasitica, 669
Bacteria: predisposition of bean plants treated with heat and inoculated with Puccinia antirrhini to bacterial infection, 733
Bacterial diseases: bacterial rhizome rot of banana, caused by Pectobacterium carotovorum (syn. Erwinia carotovora), in Jamaica, 214; common blight of bean, caused by Xanthomonas phaseoli, various bean lines show differential reaction in pods and leaves to X. phaseoli, 278; leaf scald of sugarcane, a method for isolating Xanthomonas albilineans from sugarcane leaves is described, 439; pea blast, caused by Pseudomonas syringae, in Chile, 568; bacterial leaf spot of Rieger begonia, Xanthomonas begoniae will survive for at least a year in dry leaves and be pathogenic, 814; leaf freckles and wilt of corn, resistance in 113 corn lines is tested by artificial inoculation, 956; Pseudomonas syringae is cause of a leaf necrosis of wheat in South Dakota, 1061; bacterial glume blotch of rice, caused by Pseudomonas oryzicola, is reported for first time in Australia, 1126
---: bacterial wilt of tomatoes: use of virgin soil in greenhouse for control of Pseudomonas solanacearum, 86; P. solanacearum is identified in Guam on tomato and eggplant, 793
---: fire blight of apple: inoculation trials with Erwinia amylovora, 243; smaller fruits and phytotoxicity resulted from low volume spraying of 'Jonathan' apple trees, 620; field susceptibility ratings for 46 apple cultivars, 819
Barley: Ascochyta graminea was collected from barley cultivars in Virginia, 455; barley yellow dwarf virus, mixed infections of virus isolates occurred in winter barley in New York, 472; Cephalosporium stripe was found on winter barley in Kansas in 1973, 566; male-sterile barleys from the World Collection of barley were tested for sources of resistance to ergot, 679; pathogenicity of Cylindrocladium species to 'Fla. 102' barley, 1017
---: Helminthosporium stripe: four inoculation techniques compared for the most effective method of inducing Helminthosporium stripe infection in barley, 32; a method is presented for evaluating the intensity of Helminthosporium stripe of barley, caused by H. gramineum, 265
Bean (Phaseolus vulgaris): differential reactions of pods and foliage of bean lines to Xanthomonas phaseoli suggest that different genetic systems control the reactions of pods and leaves to the bacterium, 278; bean common mosaic virus, caused yield loss in Morocco, 307; rust, caused by Uromyces phaseoli, found for first time in Wisconsin in 1973, 330; lima (Phaseolus lunatus) and snap bean (P. vulgaris) are reported as new hosts of Cristulariella pyramidalis, 421; Fusarium solani f. sp. pisi and F. solani f. sp. phaseoli were tested in combination and individually on peas and beans, 500; Fusarium equiseti and F. moniliforme found to be seedborne on bean

in Argentina, 507; predisposition of bean to bacterial infection from snapdragon rust after heat treatment, 733; resistance to Pythium ultimum found in a white-seeded snap bean line 1273, 774
---: nematodes: lima bean yields increased by control of Meloidogyne incognita with nematicides, 749; root-knot nematode control with DD and EDB increased bean yields, 1020
---: ozone: ten varieties of bean were tested for susceptibility to ozone, 14; primary leaves of 'Pinto' bean were protected from ozone injury by carboxin and other oxathiin derivatives, 162
---: root rot: only four of 125 bean cultivars and breeding lines showed promising tolerance to the bean root rot complex, 229; root rot on dryland pinto beans in Colorado decreased yields, 890; BAY 22555 at .48 lb/A was the only fungicide tested that provided significant control of root and hypocotyl rot of bean in the field in Wisconsin, 941
---: white mold: postharvest control of Sclerotinia rot of snap bean pods with heated and unheated chemical dips of DCNA and thiabendazole, 59; modifications of plant architecture can have an effect on severity of white mold infection in Great Northern dry bean varieties, 379; 'Black Turtle Soup', 'Sanilac', 'Capitol', 'Aurora', and NY 69-6207-2 edible dry beans showed tolerance to a white mold epidemic in Nebraska, 782
Beech: beech bark disease complex, caused by Nectria coccinea var. faginata, Cryptococcus fagi and Xylococculus betulae, survey in Pennsylvania shows an extension of 180 miles west and 20 miles south from previously known locations, 718
Beet (see Sugarbeet)
Begonia: Xanthomonas begoniae, isolated from leaves of Rieger begonia and stored in an envelope in a desk drawer for a year was pathogenic when inoculated into healthy leaves, 814; Schwabenland Red begonia and cultivars derived therefrom were susceptible to powdery mildew, four fungicides gave protection; Aphrodite types were immune, 875; foliar nematode, Aphelenchoides fragariae, was controlled on Rieger begonia with foliar applications of oxamyl, 911
Belonolaimus longicaudatus: 374; 822
Bermuda: 974
Beta vulgaris (see Sugarbeet)
Big bud: of tomato, found in New York in 1973, 211
Bioassay: to compare organic and copper fungicides for their biological activity, and their tenacity for control of tea leaf blister blight, 928
Biological control: discovery of Cercospora piaropi as a pathogen of water hyacinth in Florida suggests this organism as a possible source for biological control, 277; possible practicality of treating pruning wounds of apricot trees with Gliocladium roseum, which in cross-plating with Eutypa armeniacae showed typical mycoparasitic reaction, 983
Bipolaris maydis (see also Helminthosporium maydis): 529
Bispora sp.: 180
Black locust (Robinia pseudoacacia): severe dieback of young locust trees on a mine reclamation project was caused by Nectria cinnabarina, 872
Black walnut (see Walnut)
Blast: pear blast, caused by Pseudomonas syringae, was found in Chile, 568
---: rice blast, a distinct diurnal and nocturnal periodicity of conidial release was observed in the conidial release of Pyricularia oryzae over a rice blast nursery, 544
Blight: Fusarium blight of Kentucky bluegrass, variety, mowing height, and fertilization affect incidence, 514; chestnut blight, Endothia parasitica was recorded for the first time on live oak in North Carolina and Florida, 596; pod and stem blight of soybean, caused by Diaporthe phaseolorum var. sojae, was reduced by benomyl spray, 760; leaf blister blight of tea, copper fungicides appear to be more effective than organic fungicides because of their tenacity under extremely heavy rainfall, 928; spur blight of red raspberry, no immunity was found in 144 cultivars and selections, but there is a range in susceptibility, 1024
---: citrus blight (see also Young tree decline): a decline of unknown cause, affected trees have very restricted uptake of water injected into their trunks in comparison with healthy looking trees, 801; scanning electron microscope photographs show vessel-obstructing material in xylem of blight-diseased citrus trees, 1051
---: fire blight of apple: inoculation trials with Erwinia amylovora to apple rootstocks, 243; dilute and

597-490 O - 75 - 2

concentrated sprays of streptomycin compared for effect on uptake by apple trees and the effect on fruit size, 620; susceptibility ratings of 46 apple cultivars to fire blight were compiled for three North American geographic-climatic regions, 819
---: leaf, of corn, Helminthosporium turcicum and H. maydis can affect corn root and top growth at the seedling stage, 219; effect on top weight, root weight, and root rot of corn seedlings by three Helminthosporium leaf blights, 313; racial composition of isolates of Helminthosporium maydis and H. carbonum collected from corn in normal cytoplasm in Pennsylvania, 934
---: southern corn leaf blight (see Corn)
Blueberry: fungi associated with blueberry stems in Michigan, 180; triforine was superior to all other fungicides tested in control of primary infection of highbush blueberry by Monilinia vaccinii-corymbosi, 840
Blue spruce (see Picea pungens)
Botryodiplodia theobromae: 1134
Botryosphaeria ribis: 1030
Botrytis cinerea: 180; 391; 558; 915
Brassica carinata: Fusarium oxysporum f. sp. conglutinans race 1 was pathogenic to this potential vegetable crop, but race 2 was not, 479
Bremia lactucae: 118
British Columbia: 1024
Bunt: dwarf, of winter wheat, caused by Tilletia controversa, thiabendazole seed treatment gave fair to good control but caused some yield loss at 8 oz/100 lb rate, 743; seedborne bunt of wheat, caused by Tilletia caries and T. foetida, was controlled in Australia with BAY 22555 as a seed treatment, 845

Cajanus cajan (arhar): found to be susceptible host of Heterodera vigni in India, 345
Calcium nutrition: oat cultivars resistant to wilt toxin, victorin, always had higher calcium content than intermediate and susceptible types, 3
California: 54; 88; 94; 99; 118; 478; 633; 663; 869; 904; 1002
Camellia: C. sasanqua seedlings are more sensitive to dieback and canker, caused by Glomerella cingulata, than are S. japonica seedlings, 139; G. cingulata, pathogenic to camellias, was found to produce conidia in abundance by physically scraping 3- to 6-day-old cultures grown on carrot juice agar (CJA) or CJA + orange peel, 300; G. cingulata was controlled on Camellia sasanqua with benomyl-treated soil, or with spray, 710
Canada: 342; 677; 1024; 1091; 1105
Canada thistle: mycoplasmalike bodies were found in the phloem of leaf vein samples that showed typical yellows symptoms, 906
Canker: and dieback of Camellia, relative susceptibility of two species to Glomerella cingulata, 139; Eutypa canker of apricot in Spain, 442; pitch canker of slash pine, in slash pine plantations in Florida, 451; of Maesopsis trees and seedlings, Fusarium solani and Volutella sp. inoculated jointly into seedlings caused more severe cankering than either pathogen alone, 463; Nectria canker of black locust in New Brunswick, Canada, 872; Scleroderris canker, introduction and spread of disease by Christmas trees is possible, 892; wilting and dieback of stems of Rhamnus frangula 'Tallhedge' are associated with basal stem cankers caused by Tubercularia ulmea, 937; Leucostoma canker development in artificially inoculated potted peach trees, periodic water stress does not influence susceptibility of young trees to the disease, 971; of sugar maple, the time of origin of most cankers dissected was during the last 10 years and particularly the last 2 years, 1042; stem girdling canker of black walnut, is thought to be caused by symbiosis between Fusarium fungi and ambrosia beetles, 1044; Scleroderris canker on red pine nursery stock was controlled with chlorothalonil and chlorothalonil + cycloheximide, 1097
Cannabis sativa (hemp): host of Cercospora cannabina in experimental planting in Mississippi, the first record for the pathogen in North America, 165
Capital Territory: 92
Capsicum annuum (see Pepper)
Carrot: tests with Thielaviopsis basicola, if valid, confirm previous results that Thielaviopsis is one of the few microorganisms which can be pathogenic in one situation and symbiotic in another, 490; carrot motley dwarf virus and the vector for this virus were found in Oregon, 766
Cashew (Anacardium occidentale): Phytophthora nicotianae var. nicotianae is cause of shoot rot and leaf fall, 767
Cassava: root rot, caused by Phytophthora drechsleri, in Colombia, 703

Cavariella aegopodii: 766
Celery: Dyrene (anilazine) or chlorothalonil at 1.5 lb/acre gave good control of Septoria leafspot, 232
Centaurea cyanus: new host of tobacco streak virus, 354
Cephalosporium sp.: 180
--- gramineum: 566
Ceratocystis fagacearum: 65; 92; 622; 738
--- ulmi: 382; 511; 727; 784
Cercospora sp.: 332; 1015
--- arachidicola: 666
--- beticola: 434; 952
--- cannabina: first North American record for C. cannabina, found on Cannabis sativa in Oxford, Mississippi in August 1972, 165
--- cannabis: 165
--- kikuchii: 50; 130
--- oryzae: 507
--- piaropi: 277
--- zeae-maydis: 456; 532
Cercosporella herpotrichoides: 554
Certification: indexing of new strawberry indicator clones for detection of 12 viruses, 28
Chactocnema confinis: 239
Chamaecyparis lawsoniana: soil drench of ETMT before planting controlled Phytophthora cinnamomi root rot in Australia, 437
Chemicals (see also Antibiotics, Fungicides and Nematicides): 2,4-D amine was not effective against elm bark beetles in elms affected with Dutch elm disease, 382
---: cacodylic acid injected into Dutch elm-diseased trees prevented elm bark beetle development, 382
---: carbon:nitrogen ratio in soil and its relation to survival of Macrophomina phaseolina in soybean stems in soil, 1034
---: erythorbic acid, sodium erythorbate, ascorbic acid and/or sodium ascorbate soil and spray treatments contributed to recovery of orange and grapefruit trees from young tree decline symptoms, 918
---: ethylene, potato plants grown in a growth chamber without filtered air showed air pollution injury from ethylene, 304
---: herbicides, addition of preplant incorporated and preemergence herbicides (prometryn and trifluralin) to nematode-infested cotton caused additional plant injury, 120
---: lead arsenate, azinphosmethyl, sulfur 90, demeton 25, and dieldren, 247
---: methyl alkanoates and aliphatic alcohols effectively eradicated powdery mildew from dormant apple buds, 949
---: nitrogen, high nitrogen fertilizer application in fall alleviated air pollution-induced symptoms exhibited by eastern white pine Christmas trees, 150
---: oxydemeton-methyl was effective in preventing development of elm bark beetles in Dutch elm-diseased elms and in destroying broods established in trees before treatment, 382
---: sodium azide as a fungicide mixed in soil increased yields of peanuts planted in naturally Cylindrocladium black rot-infested fields, 348
---: 2,4-xylenol and m-cresol, emulsified, controlled aerial galls on apple, 701
Chenopodium amaranticolor: virus from white ash was transmitted mechanically to C. amaranticolor and identified as tobacco mosaic virus, 536
--- quinoa: successful transmission of virus isolate from white ash to C. quinoa was identified as tobacco mosaic virus, 536
Cherry: sweet cherry raspleaf virus incidence is spreading slowly but steadily in Colorado, 26; CW-524 (triforine) in greenhouse experiments exhibited after-infection activity against cherry leaf spot fungus, Coccomyces hiemalis, 326
Chickpea (Cicer arietinum): Ascochyta rabiei, a destructive pathogen of chickpea (a potential legume crop for Western Canada) was found for first time in North America at Saskatoon, Canada in 1973, 342
Chile: 568; 599
Christmas tree diseases: tip dieback of white fir, 94; air pollution symptoms were reduced on eastern white pine by fall application of high nitrogen fertilizer, 150; Phytophthora cinnamomi root rot of Fraser fir in North Carolina, 318; Gremmeniella abietina, the fungus that causes Scleroderris canker can survive in Christmas trees brought indoors, therefore, spread of fungus to other areas is possible via shipping, 892; a needlecast of blue spruce Christmas trees caused by Rhizosphaera kalkhoffii was reported for first time from North Central States, 1094
Chromatography: thin-layer chromatography used for indexing of exocortis virus-infected citron, 82
Cicadulina mbila: 968
Citrus: Phytophthora root and stem rot in milkweed vine caused by Phytophthora citrophthora isolate was not pathogenic

to citrus, 355; control of melanose on Texas grapefruit with fungicides, 504; techniques for determining susceptibility to Phytophthora parasitica, in lemon, 713; citrus nematode, Tylenchulus semipenetrans populations were sharply reduced with foliar applications of oxamyl but results were erratic, 882; occurrence of benzimidazole-resistant strains of Penicillium digitatum in Florida citrus packinghouses, 933; biphenyl, durene, and PMB were equally effective in control of green mold of artificially inoculated grapefruit, 1143
---: virus: use of thin-layer chromatography for indexing exocortis virus, 82; suppression of tristeza virus symptoms in Mexican lime seedlings by warm temperatures was a masking effect, 757; four citrus viruses, psorosis, concave gum, infectious variegation and vein enation viruses were inactivated by warm temperatures in a greenhouse for 3 to 4 months, tristeza with supplemental hot-moist air at 50°C for 3 to 7 hours was inactivated, 850
---: young tree decline (YTD): and sand hill decline, results of initial indexing tests on citrus trees to determine virus content of affected trees, 35; use of a remote sensing infrared thermometer for previsual diagnosis of YTD was not successful, 793; diagnosis of young tree decline, blight, and sand hill decline by measurement of water uptake using gravity injection, 801; three of five YTD-infected trees responded to treatment with tetracycline, 895; chemotherapy with aqueous erythorbic acid, sodium erythorbate, ascorbic acid and/or sodium ascorbate resulted in extensive growth of new foliage and substantial recovery of YTD trees from the disease symptoms, 918; xylem of blight-diseased citrus trees was photographed with a scanning electron microscope which revealed vessel-obstructing material, 1051
Cladosporium sp.: 208; 332
Claviceps purpurea: 679
Clover (white): peanut stunt virus found on white clover in Florida in two widely separated counties, 830
Coccomyces hiemalis: 326
Coconut: a new record of infection of old coconut palms by Ganoderma boninense in Sri Lanka, 293
Collecephalus hemerocalli: 102
Colletotrichum gloeosporioides: 1134
--- graminicola: 532
--- pisi: 226
Colocasia esculenta (dasheen): was found infected with dasheen mosaic virus in Venezuela, 1032
Colombia: 703
Colorado: 26; 110; 571; 890
Coniothyrium sp.: 180
Connecticut: 40; 428
Conoderus falli: 239
Corn: a new virus disease found in Venezuela is not mechanically transmissible and the vector is a leafhopper, Peregrinus maidis, 122; two biotypes of Meloidogyne hapla reproduced on corn in the Pacific Northwest trials, 128; corn was successful as fallow in reducing the nematode population of Rotylenchulus reniformis in a sweetpotato nematode control program, 259; stalk rot, 50 corn fields surveyed in Minnesota for stalk rot and lodging, 363; occurrence of Phomopsis on maize (corn), 416; gray leaf spot, caused by Cercospora zeae-maydis, is associated with no-tillage production of corn, 456; Colletotrichum graminicola, Puccinia polysora, and Cercospora zeae-maydis were among the most damaging pathogens observed on corn in North Carolina in 1972 and 1973, 532; symptoms induced in corn by Exserohilum prolatum, a recently discovered pathogen, were generally similar to those induced by E. rostratum (=Helminthosporium rostratum), 612; seed- and soilborne fungal complex, the cause of severe stand losses in 'Florida Sweet' corn, was curbed with a seed slurry treatment of benomyl and captafol (each at 2 oz per cwt), 922; 113 corn lines were tested for reaction to leaf freckles and wilt caused by Corynebacterium nebraskense, 956; "red kernel" of sweet corn on the market was caused by Epicoccum nigrum, 1065; detection of spiroplasma in corn stunt-infected vector, Dalbulus maidis, supports the hypothesis that the CS spiroplasma is the corn stunt disease agent, 1109; effect of cob and shank inoculations with Diplodia maydis on cell death in stalk internodes of corn, 1113
---: downy mildew: downy mildew, caused by Sclerospora sorghi, was found on corn in Kentucky, 90; sorghum downy mildew on corn in India is being perpetuated by a collateral host, Heteropogon contortus, 285; first report of its occurrence on corn in Indiana, 430; inoculation technique described for inoculating corn and sorghum with

Sclerospora sorghi to assess resistance, 825
---: leaf spot: 76 corn inbred lines varied in reaction to Helminthosporium leaf spot caused by isolate 72: 44-6 (ATCC 24962), 399; field reactions of corn inbreds to a Helminthosporium leaf spot caused by isolate 72: 44-6, a Helminthosporium resembling race 2 of H. carbonum in pathogenicity pattern, but differing from race 1 and 3, 909; seedling reactions of 87 corn hybrids to Helminthosporium leaf spot isolate 72: 44-6, 975
---: southern corn leaf blight: fungicide treatments with Dyrene, zinc + maneb, chlorothalonil and copper salts of fatty and rosin acids (C 4-E) reduced severity of southern corn leaf blight on corn containing Texas male-sterile cytoplasm and increased yields, 104; root and top growth at seedling stage is affected by Helminthosporium blights and is measurable, 219; Helminthosporium maydis race T sporulation with virtually none at 93% relative humidity (RH), increased with increasing RH so that as many spores were formed at 97-98% RH as were previously reported found when free water was present, 297; leaf infections of H. turcicum and H. maydis races O and T reduced top and root weights of corn seedlings and resulted in more root rot infection by Fusarium moniliforme, 313; maintenance of stock cultures of H. maydis (races O and T), 334; Bipolaris maydis (H. maydis) race and mating type frequencies have changed in North Carolina, 529; presence of "Texas" cytoplasm in corn varieties detected with a simple biochemical assay by using H. maydis race T pathotoxin, 777; reactions in corn inbreds and hybrids to H. maydis race T were determined by measuring lesion sizes on adult plants inoculated in field plots, 811; racial composition of H. maydis and H. carbonum on corn hybrids in normal cytoplasm in Pennsylvania in 1973, 934
Correction: (Vol. 51, No. 11, Page 936), 192
---: (Vol. 57, No. 11, Pages 960-963), 576
---: (Vol. 58, No. 2, Page 171), 288
---: (Vol. 58, No. 4, Page 306), 768
---: (Vol. 58, No. 8, Page 683), 1056
---: (Vol. 58, No. 8, Page 730), 1056
---: (Vol. 58, No. 9, Page 791), 960
Corynebacterium nebraskense: 956
Corynespora cassiicola: 636
Coryneum microstictum: 180
Cosmos bipinnatus: new host of tobacco streak virus, 354
Cotton: soil containing 9.5 ppm of carboxin protected potted plants against ozone, 162; a first report of Alternaria macrospora attacking cotton (Gossypium hirsutum) in the Southeastern United States, in Louisiana in 1972, 201; low cotton and soybean yields in east Georgia may be caused by hardpan, low pH, and nutritional factors as well as by nematodes, 369; reduction in yield of cotton caused by parasitic diseases in 1973, 373; subsoiling under the planting row (SUR) reduced symptoms of cotton stunt disease complex by increased root development, shoot growth and yields of three cotton cultivars in Georgia, 541; first report of Verticillium dahliae on cotton in Iraq, 996
---: nematodes: cotton seedlings under stress from root-knot nematode infection were susceptible to more injury by addition of herbicides, 120; chemical and cultural control of reniform nematode with nematicides and crop rotation, 337
Cowpea (Vigna unguiculata ssp. unguiculata): infection by uredospore suspensions from rusted snapdragons following heat treatment, 733; an aphid-borne virus disease of irrigated cowpea, 1080
---: nematodes: Vigna spp. are hosts of Heterodera vigni in India, 345; inheritance of resistance in cultivars Mississippi Silver and Victor K798 to root-knot nematode is apparently governed by a single dominant factor, susceptibility level is under polygenic control, 361; combination treatment of 1,3-D and ultra-high frequency electromagnetic energy in soil gave excellent control of reniform nematode and common purslane, 985; an Egyptian population of Heterodera cajani caused reduction in growth and other malformations in cowpea, 1130
Crambe: (C. abyssinica and C. hispida): Fusarium oxysporum f. sp. conglutinans race 2 was pathogenic to Crambe but race 1 was not, 479
Criconemoides spp.: 169; 991; 1048
--- ornatus: 822
--- xenoplax: 76; 633
Cristulariella pyramidalis: 421
Cronartium comandrae: 616
--- fusiforme: 198; 497; 584; 1137
Crop rotation: for control of reniform nematode in cotton fields, rotation or sorghum-cotton reduced pretreatment nematode counts, 337
Crop sequence: and fallow, its effect on

populations of Rotylenchulus reniformis in fumigated and untreated soil, 259
Cryptococcus fagi: 718
Cucumber (see Cucurbits)
Cucurbits: Fusarium wilt of muskmelons found in Washington, 10; two accessions of Cucumis metuliferus are highly resistant to watermelon mosaic virus-1 and hypersensitive resistant to squash mosaic virus, 735
---: cantaloup: yield and fruit grade increased by control of Meloidogyne incognita, 746; PI 321005 (Tainan #2) from Taiwan showed high level of resistance to powdery mildew, downy mildew and gummy stem blight in Georgia tests, 899
---: cucumber: Thielaviopsis basicola may increase plant growth, 490; bacterial soft rot, Alternaria rot, black rot and Fusarium were the major causes of cucumber fruit decay besides mechanical injury and chilling in seven New York retail stores, 558; target leafspot, soil rot and powdery mildew controlled on cucumber with benomyl, sodium azide, and chlorothalonil and zinc + maneb, respectively, 636
---: watermelon: several fungicides provided some control of damping-off and seedling wilt, caused by Fusarium spp. and Rhizoctonia spp., 114; 'Charleston Gray', 'Garrisonian', and 'Florida Giant' were tested for the influence of watermelon mosaic virus on their growth habits, fruit yield and size, 195
---: virus: watermelon mosaic virus (WMV), WMV-2 caused severe damage to cucurbits in Morocco, 143; WMV-2 was found widely distributed in central valleys of Chile on squash, honeydew melon and zucchini squash, 599
Culture medium: antagonism between ungerminated spores of Trichoderma viride and Stereum purpureum, 71; attempts to support growth of the mycoplasmalike organism agents from diseased aster yellows and corn stunt plants and leafhoppers was unsuccessful, 797
Cuscuta sp. (dodder): on petunia and dusty miller, found for first time in Bermuda, 974
Cylindrocarpon sp.: 180
--- tonkinense: 1134
Cylindrocladium crotalariae: 348; 1017
--- floridanum: 705; 1017
--- scoparium: 188; 1017
Cytoplasm (male-sterile): fungicidal control of southern corn leaf blight on Texas male-sterile corn, 104; effect of relative humidity on sporulation by Helminthosporium maydis on corn, 297; reduced top and root weights of corn seedlings infected with races O and T of H. maydis, 313; detection of "Texas" cytoplasm in corn varieties by a biochemical assay by using race T pathotoxin of H. maydis, 777; differences in lesion sizes among cms-T corn plants inoculated with H. maydis race T are due to the different nuclear genes interacting with the cytoplasm, 811
Cytospora: 388
Cytosporina (Eutypa armeniacae): 869

Dactylium dendroides: 987
Dalbulus maidis: 797; 1109
Damping-off: of watermelon, soil fungicides provide some control, 114
Dasheen: infection with mosaic virus in Venezuela, 1032
Daylily (Hemerocallis): a leaf-streak disease, caused by Collecephalus hemerocalli found in Pennsylvania, 102
Decline: oak decline of scarlet oak in Virginia, large-diameter trees declined and died faster than small-diameter trees, 396; of ohia trees in Hawaii, the distribution and possible role of Phytophthora cinnamomi in the decline, 1069
Delaware: 443
Delayed harvest: of soybean in Indiana resulted in increase of seed infection and poorer quality seed, 130
Dendrophoma sp.: 180
Diaporthe medusaea: 504
--- phaseolorum var. sojae: 50; 130; 173; 411; 760; 978
--- (Phomopsis): 332
--- --- vaccinii: 180
Didymella applanata: 1024
Dieback: of Camellia, caused by Glomerella cingulata, C. sasanqua is more susceptible than C. japonica, 139; benomyl-treated soil controlled G. cingulata, 710; tip dieback of juniper in Wisconsin is associated with Sclerophoma pythiophila, 653
Dieffenbachia picta: control of Phytophthora palmivora with pyroxychlor, 1100
Diplodia zeae: 313; 414; 1113
Dips: root-dip of onion bulb with thiophanate-methyl or benomyl gave excellent control of white rot of onion in Washington, 6; postharvest control of Sclerotinia rot of snap bean pods with DCNA and thiabendazole, 59
Discosia pini: 677
Disease control: horizontal and vertical resistance discussed in relation to variety development, using Lycopersicon : Fusarium host : pathogen as

model, 579
Disease ratings: of corn hybrids and inbred lines inoculated with two isolates of Exserohilum prolatum, a recently discovered pathogen, 612
Disease resistance: selected poplar clones were evaluated for resistance to poplar leaf rust, 721
Diseases: serious and unusual plant diseases found in Connecticut in 1973, 428; of sunflower in Tanzania, 896
Distribution: distribution of beech bark disease organisms, Nectria coccinea, Cryptococcus fagi and Xylococculus betulae, in northeastern Pennsylvania, 718; of nematodes in soil from Coastal Plain, Piedmont and Mountains of North Carolina, 991
Dodder: found for the first time in Bermuda on petunia, 974
Dogwood: infection with Armillaria mellea, 719
Dollar spot: on Kentucky bluegrass, caused by Sclerotinia homeocarpa, incidence varies with variety, mowing height and fertilization level, 514
Downy mildew: of head lettuce, found in California, 118; of pea, first epiphytotic of this disease in Wisconsin, 226; of cantaloup, resistance to Pseudoperonospora cubensis in Taiwan introduction PI 321005, 899
---: Sclerospora sorghi: S. sorghi found in Kentucky on sudangrass, sudan-sorghum hybrids, corn and on shattercane (Sorghum bicolor) 90; S. sorghi on corn in India is becoming a limiting factor in production; Heteropogon contortus is collateral host, 285; on corn, this is first report of its occurrence in the Corn Belt States north of the Ohio River, 430; use of sporulation-ready donors of either corn or sorghum plants for a quantitative measurement of conidial inoculum of S. sorghi for seedlings, 825
Drechslera sorokiniana: 507
Dutch elm disease (DED): DED moved into Oregon from Idaho, 75; soil injections around large nursery elms with benomyl greatly reduced symptom expression of DED and reduced mortality, 261; oxydemeton methyl and cacodylic acid pressure-injected into sap stream of diseased elm gave complete control of elm bark beetles, 382; colonization by Ceratocystis ulmi in benomyl-treated elms was inhibited, 511; present distribution of DED in Colorado, 571; response of elm interspecific hybrids to controlled inoculation with C. ulmi, 727; 16 American elm trees from a collection of over 21,000 seedlings survived systemic infections induced by inoculation of C. ulmi into mainstems but resistance was poorly transmitted to seedling or clonal progenies, 784; treatment and inoculation of American elm trees with C. ulmi and the antibiotic nystatin the same day resulted in significant control of the disease, but not if treatment preceded or followed inoculation by a week, 924
Dwarf bunt: effect of thiabendazole on incidence of dwarf bunt and yield of winter wheat, 743
Dying arm disease: of grape, Cytosporina, the imperfect stage of Eutypa armeniacae has been consistently isolated from diseased tissues of dying arms, a newly recognized disease in California, 869

Early dying disease: of potato in Florida, possibly a Verticillium wilt-nematode complex is involved, 374
Eggplant: bacterial wilt in Guam is caused by Pseudomonas solanacearum, 793
Egypt: 176; 1130
Eichhornia crassipes (see Water hyacinth)
Electrolyte loss: electrolyte leakage from resistant and susceptible oat cultivars treated with victorin, 3
Electron microscopy: revealed viruslike particles in leaves from diseased Eucalyptus trees, 92; of new corn virus from Venezuela, 122; of artichoke latent virus, 691; rickettsia-like bacteria typical of Pierce's disease were found in grape vines at College Station, Texas, by means of electron microscopy, 780; scanning electron microscope aspects of blight disease of citrus, 1051
Elm: Dutch elm disease (DED): 75; 261; 382; 511; 571; 727; 784; 924
---: elm phloem necrosis, found for the first time in New Jersey at Phillipsburg, in 1973, 387
Endogone spp.: 169
--- mosseae: mycorrhizal development in soybean was not affected by the presence of Pythium or Phytophthora in soybean roots, 221
Endomycorrhizal fungi: extent of development of endomycorrhizal fungi on soybean roots in Florida appeared to be affected by nematode incidence, 169
Endothia parasitica: 596
Epicoccum sp.: 180
--- nigrum: 1065
Epidemic: Naemacyclus needlecast of

Scots pine in Pennsylvania, 287; several dry edible bean varieties showed tolerance to a natural epidemic of white mold disease in Nebraska, 782
Epidemiology: a study was made of the dispersal of the conidia of Pyricularia oryzae and its relationship to time and weather, 544
Epiphytotic: of anthracnose and downy mildew of pea in Wisconsin, 226; of sugarbeets, in California, caused by Erysiphe polygoni, 904
Eradication: of potato wart from West Virginia, 291
Ergot: of barley, susceptibility of male-sterile barleys, 679
Erwinia amylovora: 181; 243; 819
--- carotovora: 214; 558
Erysiphe cichoracearum: 636
Escarole: Bidens mottle virus was responsible for the majority of infections in commercial escarole fields in Florida, 1087
Eucalyptus macrorhyncha: viruslike particles isolated from diseased trees in Australia, 92
--- marginata: control of Phytophthora cinnamomi root rot with soil drench of ETMT 24 hours before planting, 437
Eutypa armeniacae: 442; 869; antagonism with Gliocladium roseum, 983
Exobasidium vexans: 928
Exserohilum prolatum: 612

Fertilization: high spring fertilization of bluegrass increased incidence of Fusarium wilt in Kentucky bluegrass cultivars, 514
Festuca rubra: new host species of Puccinia pygmaea, 1028
Fig (Ficus carica): phenamiphos applied as a paste to stem and trunk controlled root-knot nematodes on fig plants, 1075
Fire blight (see Blight)
Flax: a new and dangerous race of flax rust (race 370) was discovered in flax-producing area of North America, 311; Drechslera sorokiniana and Fusarium moniliforme were found on flax seed in Argentina, a new record, 507
Florida: 35; 165; 277; 374; 445; 451; 490; 596; 669; 763; 801; 830; 878; 895; 918; 922; 933; 1087
Foliar diseases (see also Leaf diseases): of corn in North Carolina in 1972 and 1973, 532
Fomes annosus: 282; 409; 478; 1077
--- applanatus: 110
--- lignosus: 295
Forecasting: disease forecasting method for Cercospora leaf spot of peanut developed by Jensen and Boyle aided effective control with judicious use of fungicides, 666
Fraxinus americana (white ash): tobacco mosaic virus is causal agent of a mosaic disease found in ash trees in Massachusetts, 536
Freezing stress: 'Tallhedge' (Rhamnus frangula) is apparently predisposed to attack by Tubercularia ulmea canker infection by freezing stress, 937
Fumigation (see Antibiotics, Chemicals, Fungicides or Nematicides)
Fungi: pathogenic and endomycorrhizal fungi associated with soybean roots in Florida, 169; 19 species of fungi identified from section of woody tissue from blueberry plantings in Michigan, 180; pathogens present in seed samples of crops from Argentina are recorded, 507; survey of Verticillium malthousei, Mycogone perniciosa and Dactylium dendroides in commercial mushroom houses, 987
Fungicides (see also Antibiotics, Chemicals and Nematicides): rates of pesticides commonly suggested for 100 gallons of "dilute" spray gave economic control of diseases, insects and mites on apples during 1973 when applied on an acre basis; in general, control did not increase with concentration of spray materials, 247
---: several tested together: DCNA, benomyl, thiophanate-methyl and dichlozoline effectively controlled onion white rot when dusted into the planting furrow, 6; for control of powdery mildew of Antirrhinum, 12; for control of watermelon damping-off and seedling wilt, 114; six systemic fungicides were tested against peach brown rot and scab, 208; for control of Cylindrocladium black rot, none of fungicides tested was effective, 348; for control of "peach tree decline," postpruning sprays were not effective, 388; for postharvest fruit rot of red raspberry, 391; a combination of benomyl and PCNB gave good control for stripe smut and melting out of 'Merion' Kentucky bluegrass, 573; triforine was superior to all other fungicides tested for primary control of Monilinia vaccinii-corymbosi on highbush blueberry, 840; for control of Cercospora leaf spot of sugarbeet, 952; benomyl was best of those tested for control of Phomopsis blight on juniper, 1012; for control of Cercospora leaf spot of Ligustrum japonicum, 1015;

for control of Scleroderris canker of red pine, 1097; for control of Phytophthora palmivora on Dieffenbachia picta, 1100
---: anilazine, was not an effective substitute for PCNB in control of Helminthosporium vagans, 573
---: BAS 3201 F, was effective against peach brown rot and scab, 208
---: BAS 3460 F, used for preharvest fungicide treatment for red raspberry, 391
---: BAS 67054 F, controlled powdery mildew of Antirrhinum in greenhouse test, 12
---: Bay Dam 18654: Bay Dam 18654 + PCNB was effective against stripe smut and melting out in Kentucky bluegrass, 573; gave some control of mummy berry of highbush blackberry, 840
---: BAY 22555: did not control Phytophthora cinnamomi root rot of Chamaecyparis lawsoniana and Eucalyptus marginata, 437; gave outstanding control of seedborne bunt of wheat in New South Wales, 845; was the best of four fungicides tested for control of bean root rot complex in Wisconsin, 941
---: benomyl: controlled onion white rot by dusting planting furrow and by root-dip treatment, 6; reduced injury caused by ozone on annual bluegrass, Poa annua, 41; red oak seedlings treated with benomyl drench before inoculation with oak wilt fungus did not develop symptoms of wilt and pathogen was not recovered, 65; showed promise for control of watermelon damping-off and seedling wilt, 114; oxidant injury on azalea plants was suppressed by benomyl drenches, 136; benomyl and TCMTB-60W were compared for effectiveness in control of seedling and pod rot disease of peanut, 176; was effective against peach brown rot and scab, 208; benomyl and thiabendazole tolerance found in Fusarium oxysporum f. sp. gladioli on a California and a Florida farm where gladiolus corms were dipped in benomyl during 2-3 crop seasons, 256; injected into soil was effective in disease control of Dutch elm disease of large nursery elms, 261; was used as preharvest treatment for red raspberry, 391; strains of Cercospora beticola are resistant to benomyl in Texas, 434; controlled Phomopsis blight on 2-year-old nursery seedlings of eastern redcedar, 476; benomyl spray applications, postbloom and 1 month later, controlled melanose on grapefruit, 504; Dutch elm disease-inoculated elms treated with benomyl showed restricted colonization and localized disease symptoms, 511; used as a protective spray on wheat against Cercosporella foot rot, 554; gave excellent control of target leaf spot of cucumber, 636; survey for benomyl-tolerant isolates of Monilinia fructicola and M. laxa in stone fruit orchards of California yielded none on 37 isolates tested, 663; benomyl applied on a meteorological schedule by using Jensen and Boyle's forecasting system gave good control of Cercospora leaf spot of peanut, 666; gave good control of Fusicladium effusum, 687; controlled Camellia dieback, 710; as foliar applications, reduced the internally seedborne fungi in soybean, 760; bottom rot of lettuce was significantly reduced by benomyl spray, 837; gave control against mummy berry of highbush blueberry, 840; apple scab fungus showed tolerance to benomyl sprays in a 'Jonathan' orchard in Australia, 886; use of benomyl-malt agar for control of contaminants in cultures of wood decay fungi, 902; benomyl mixed with captafol as seed slurry treatment significantly improved plant stand of 'Florida Sweet' corn by controlling seed- and soilborne fungi, 922; benomyl-resistant strains of Penicillium digitatum, 933; not effective in control of bean root rot complex, 941; controlled Cercospora leaf spot of sugarbeet, 952; benomyl sprays gave good control of Phomopsis blight on juniper, 1012; controlled Cercospora leaf spot of Ligustrum, 1015
---: biphenyl, two aromatic hydrocarbons were compared for control of Penicillium digitatum-inoculated grapefruit and the three were equally effective, 1143
---: BT (RH-124), tests indicate that leaf rust control by a single application of BT can substantially increase forage yields from winter wheat, 272
---: captafol: single postbloom application controlled melanose on Texas grapefruit, 504; captafol and benomyl mixed together as seed slurry, controlled seed- and soilborne complex of diseases and improved plant stand of new corn variety, Florida Sweet, 922; gave promising results in control of bean root complex in greenhouse, but not in field tests, 941
---: captan: improved germination of soybean seed grown in Illinois, 50; used in tests for minimum rates of pesticides on apples, 247; used as preharvest treatment for red raspberry, 391;

597-490 O - 75 - 3

Fungicides (continued): captan and zinc + maneb were not effective in control of cedar-apple rust in greenhouse tests, 587
---: carboxin: carboxin and other 1,4-oxathiin derivatives protected beans against injury when exposed to ozone at 25 pphm for two hours, 162; carboxin was compared with benomyl and TCMTB-60W for control of peanut seedling and pod rot disease, 176; reduced injury from oxidants in azalea plantings, but not as effective as benomyl or thiophanate, 136
---: chloroneb: chloroneb incorporated into soil controlled damping-off of watermelon seedlings caused by Rhizoctonia, 114; was not an effective substitute for PCNB against Helminthosporium leaf spot on Kentucky bluegrass, 573
---: chlorothalonil: fungicidal control of southern corn leaf blight on Texas male-sterile corn with chlorothalonil, 104; controlled Septoria apiicola on celery, adjuvants did not improve performance, 232; as a preharvest fungicide treatment for red raspberry, 391; was effective in control of new leaf disease on Aechmea fasciata, caused by Helminthosporium rostratum, 445; as single and double applications, chlorothalonil did not control melanose, 504; gave satisfactory control of powdery mildew of cucumber, but not target leaf spot, 636; application of chlorothalonil by the Cercospora leaf spot disease forecasting method was effective in control of the disease, 666; applied to underside of lower leaves significantly reduced infection by Rhizoctonia solani on lettuce, 837; chlorothalonil and chlorothalonil + cycloheximide were effective against Lophodermium pinastri on Scotch pine Christmas trees, 853; not effective in control of powdery mildew on Rieger begonia, 875; effective against Cercospora sp. on Ligustrum, 1015; chlorothalonil and chlorothalonil + cycloheximide were best of six fungicides tested for control of Scleroderris canker on red pine nursery stock, 1097
---: copper fungicides were more effective than organic fungicides in control of tea leaf blister blight in laboratory and greenhouse studies, 928
---: copper ammonium carbonate as single or double applications controlled melanose of grapefruit, 504
---: copper salts of fatty and rosin acids (C 4-E), applications of fungicides controlled southern corn leaf blight on Texas male-sterile corn and resulted in increased yields, 104
---: tribasic copper sulfate: sprays made postbloom and 1 month later gave control of melanose on grapefruit, 504; controlled Cercospora leaf spot on Ligustrum, 1015
---: copper-sulfur dust, used in testing the disease forecasting method for applying fungicides for control of Cercospora leaf spot of peanut, 666
---: cupric hydroxide, was harmful to Concord grape yields by affecting cluster and berry weights, 486; two spray applications controlled melanose on grapefruit, 504; applied according to forecasting method for Cercospora leaf spot control on peanut, cupric hydroxide was effective, 666
---: DCNA: controlled onion white rot when dusted in furrow in Washington, 6; controlled white mold of snap bean pods during storage, 59
---: DD-MENCS, as fumigant was very effective against damping-off and seedling wilt of watermelon, 114
---: dichlozoline: as a foliar spray or as a dust in planting furrow controlled white rot of onion, 6; was effective against brown rot of peach, but not scab, 208
---: dinocap: was used in tests for minimum rates of pesticides, 247; effective in control of powdery mildew on susceptible Rieger begonia cultivars, 875
---: durene, an aromatic hydrocarbon, proved to be as effective as biphenyl in control of green mold on grapefruit, 1143
---: Dyrene (anilazine): reduced severity of corn leaf blight on corn containing male-sterile cytoplasm, 104; gave good control of Septoria leafspot of celery at 1.5 lb/acre, 232
---: ETMT: gave good disease control for a serious new crown rot and wilt of baby's breath, 669; as a preplant drench, controlled Phytophthora cinnamomi root rot of Chamaecyparis lawsoniana and Eucalyptus marginata, 437
---: ferbam, the standard registered fungicide for control of mummy berry of highbush blueberry failed to reduce primary or secondary infections or to increase yield, 840
---: HOE 2873: significantly reduced powdery mildew on Antirrhinum, 12; was a highly effective fungicide against rusty spot of peach, 254
---: maneb, provided control of needlecast of Scotch pine Christmas trees, 853
---: methyl bromide, fumigation of wheat

fields in Florida infested with soilborne wheat mosaic increased yields of susceptible cultivars, reduced yields of resistant varieties, and did not eliminate vector from soil, 878
---: NIA 9102, controlled Cercospora leaf spot on Ligustrum, 1015
---: PCNB: PCNB-ETMT provided some control of watermelon damping-off and seedling wilt, 114; PCNB in combination with benomyl was effective in control of stripe smut and Helminthosporium leaf spot of Kentucky bluegrass, 573; PCNB-BAY 22555, showed promise for control of bean root rot complex, 941
---: PMB (pentamethylbenzene), and durene were as effective as biphenyl in control of green mold of artificially inoculated grapefruit, 1143
---: pyroxychlor, gave superior control of Phytophthora palmivora on Dieffenbachia picta, 1100
---: sodium azide, appeared promising for control of soil rot of cucumber, 636
---: sulfur dusts and sprays were effective against powdery mildew of mango, 45; sulfur-zineb, used for control of Cercospora leaf spot of peanut, 666; sulfur, and sulfur with cupric hydroxide gave protection to Rieger begonia cultivars that were susceptible to powdery mildew, 875
---: TCMTB-60W (2-(thiocyanomethylthio) benzothiazole), a new systemic liquid fungicide was effective in control of seedling and pod rot disease of peanut, 176
---: thiabendazole: was effective as heated or unheated dip for control of Sclerotinia rot of snap bean, 59; gave some control of Fusarium spp. that caused damping-off and seedling wilt of watermelon, 114; used as preharvest fungicide for red raspberry, 391; was not effective against stripe smut on bluegrass, 573; as a seed treatment gave fair to good control of dwarf bunt on winter wheat, 743; thiabendazole-resistant and benomyl-resistant strains of Penicillium digitatum occur in Florida packinghouses and could cause a serious decay-control problem, 933; controlled Cercospora leaf spot of sugar beet, 952
---: thiophanate: reduced ozone injury on Poa annua, 41; suppressed oxidant injury on azalea plants, 136; tested as preharvest treatment for red raspberry, 391
---: thiophanate-methyl: root-dip and furrow applications controlled white rot of onion, 6; controlled powdery mildew on Antirrhinum, 12; controlled peach brown rot and scab, 208; was effective in control of Phomopsis juniperovora on eastern redcedar seedlings, 476; was not effective for control of Ustilago striiformis on bluegrass, 573; gave some control of mummy berry of highbush blueberry, 840; gave poor control of powdery mildew on Rieger begonia, 875; apple scab fungus showed tolerance to thiophanate-methyl and benomyl in Australia tests, 886; controlled Cercospora leaf spot of sugarbeet, 952
---: TPTH: controlled Cercospora leaf spot of sugarbeet, 952; controlled Cercospora leaf spot on Ligustrum japonicum, 1015
---: triarimol: controlled powdery mildew (Oidium sp.) on snapdragon, 12; effective against brown rot of peach, not effective for scab, 208; controlled stripe smut on bluegrass, but commercial development has been discontinued, 573
---: (triforine) CW-524: controlled brown rot of peach, 208; showed post-infection activity in control of cherry leaf spot in greenhouse experiments, 326; gave good control of cedar-apple rust on apple, 587; was superior to all other fungicides tested in controlling primary infection of highbush blueberry by Monilinia vaccinii-corymbosi, 840; gave protection to susceptible cultivars of Rieger begonia to powdery mildew, 875
---: triphenyltin chloride, controlled Cercospora leaf spot of sugarbeet, 952
---: U32104 (2-(methoxycarbamoyl) benzimidazole), + PCNB controlled stripe smut and Helminthosporium vagans in Kentucky bluegrass, 573
---: zinc + maneb: reduced infection by Helminthosporium maydis on corn in Texas male-sterile cytoplasm, 104; used to test for minimum rates of pesticides on apples, 247; controlled Helminthosporium rostratum infection on Aechmea fasciata, 445; increased wheat yields in test plots by controlling leaf and stem rust of wheat, 469; effective in control of powdery mildew of cucumber, but not for target leaf spot, 636; controlled Cercospora leaf spot of sugarbeet, 952; gave good control of leaf spot of Ligustrum japonicum caused by Cercospora sp., 1015
Fusarium sp.: 184; 332; 558; 1044
--- spp.: 114; 169; 180; 373; 896
--- lateritium f. pini: 451
--- moniliforme: 313; 363; 507
--- nivale: 432
--- oxysporum f. sp. conglutinans: 479

Fusarium oxysporum f. sp. gladioli: 256
--- --- f. sp. lycopersici: 320; 579
--- --- f. melonis: 10
--- --- f. sp. pisi: 62; 500
--- --- f. sp. vasinfectum: 373
--- roseum: 514
--- --- 'Graminearum': 363
--- solani: 229; 463; 754
--- --- f. sp. phaseoli: 500; 890; 941
--- tricinctum: 514
Fusicladium effusum: 687

Galinsoga parviflora: new host of Whetzelinia sclerotiorum, 700
Galling: of apple, aerial galls of unknown cause were controlled with 2,4-xylenol and m-cresol, 701
Ganoderma boninense: 293
Gardenia: stem and trunk applications of phenamiphos (paste) significantly reduced Meloidogyne arenaria populations in greenhouse and screenhouse experiments, 1075
Geographic distribution: of Oidium mangiferae, the causal organism of powdery mildew of mango, 45
Georgia: 21; 76; 107; 369; 553; 723
Germination: captan seed treatment improved germination of soybean seed, from Illinois, 50; low rates of germination among soybean seed lots from three regions in Illinois were associated with big percentages of Diaporthe-infected seeds, 411; Diplodia zeae and β-Phomopsis infection of corn seeds lowered the germination in blotter seed health test, 414
Gladiolus: method described for rapid assay of soils for gladiolus Fusarium corm rot inoculum potential in field samples, 184; tolerance of F. oxysporum f. sp. gladioli to benzimidazole fungicides resulted in severe corm rot when benomyl-dipped corms were planted, 256; a photocopy system was developed to facilitate the rapid and accurate evaluation of fluoride injury of gladiolus indicator plants in Ontario, 1091
Gliocladium roseum: in crossplating with Eutypa armeniacae, the pathogen that causes canker of apricot, G. roseum showed a mycoparasitic reaction and this suggests the practical application of treating pruning wounds and diseased trees with G. roseum, 983
Gloeocercospora sorghi: 267
Glomerella cingulata: 139; 300; 710
Glume blotch: of rice, caused by Pseudomonas oryzicola, is reported for the first time from Australia, 1126
---: of winter wheat in Georgia, caused by Septoria nodorum, 21
Glycine spp. (see also Soybean): species other than G. max were tested to determine whether resistance to bean pod mottle virus occurs within the genus, 191
Godronia cassandrae f. vaccinii: 180
Golf greens: nematodes associated with bentgrass and bermudagrass golf greens in North Carolina, 822
Goss's wilt (see Leaf freckles)
Grand fir (see Abies)
Grape (Vitis spp.): Cristulariella pyramidalis was found on grape in West Virginia, 421; cupric hydroxide sprays on Concord grape reduced yields, induced defoliation, and lowered juice quality, 486; deep placement and high doses of 1,3-D coupled with low soil moisture gave promising nematode control in replant vineyards, 562; Pierce's disease, thought to be the cause of malady in six grape clones at College Station, Texas; rickettsia-like bacteria found in xylem cells, 780; dying arm disease, Eutypa armeniacae, the apricot dieback pathogen appears to be closely associated with this newly recognized disease in California, 869
Grasses: bentgrass, samples from bentgrass golf greens indicated Trichodorus christiei, Hoplolaimus galeatus, Tylenchorhynchus claytoni, and Helicotylenchus dihystera may be the most important nematodes on bentgrass in North Carolina, 822
---: bermudagrass, Criconemoides ornatus, Helicotylenchus dihystera, Trichodorus christiei, Meloidogyne sp., Tylenchorhynchus claytoni, Hoplolaimus galeatus, and Belonolaimus longicaudatus were recovered from bermudagrass-over-seeded golf greens in North Carolina, 822
---: bluegrass: several cultivars were evaluated for midsummer Helminthosporium sorokinianum leaf spot, 448; incidence of Fusarium blight and Sclerotinia dollar spot on bluegrass cultivars varied with variety, mowing height and fertilization level, 514; a single fall application of benomyl and PCNB combination gave excellent control of stripe smut and leaf spot on 'Merion' bluegrass, 573; hyphae of Sclerophthora macrospora were found consistently associated with bluegrass and bentgrass suffering from yellow tuft disease, 848
---: johnsongrass: is host of maize chlorotic dwarf virus in Illinois, 420
---: Poa annua, ozone injury reduced by

systemic fungicides, 41
---: red fescue: disease reaction of 56 introductions and 5 cultivars to Puccinia pygmaea, 1028
---: Rhodesgrass (Chloris gayana), found infected by Helminthosporium hawaiiense; first report of pathogen from continental United States, in Florida, 490
---: shattercane (Sorghum bicolor), first report of downy mildew on shattercane in Kentucky, 90
---: sudangrass and sudan-sorghum hybrids, occurrence of downy mildew, caused by Sclerospora sorghi, in Kentucky, 90
Green mold: of grapefruit, biphenyl compared with durene and PMB for control, 1143
Gremmeniella abietina (Scleroderris lagerbergii): 892
Growth loss: of scarlet oak, a 50% growth loss occurred in oak decline-affected trees for the period following decline initiation until tree death, 396
Guam: 793
Gummy stem blight: of cantaloup, caused by Mycosphaerella citrullina, resistance to this pathogen as well as Sphaerotheca fuliginea race 2 and Pseudoperonospora cubensis was shown in cantaloup introduction from Taiwan, 899
Gymnosporangium clavipes: 696
--- juniperi-virginianae: 587; 696

"Hardcore": of sweetpotato, flexuous rod particles were isolated from foliage of sweetpotato plants produced from plants having "hardcore," 17
Hawaii: 102; 401; 495; 1069
Helicotylenchus sp.: 991
--- spp.: 1048
--- dihystera: 169; 822
Helminthosporium (72: 44-6): 399; 909; 975
--- avenae: 80
--- carbonum: 934
--- gramineum: 32; 265
--- hawaiiense: 490
--- maydis (see also Bipolaris maydis): 104; 219; 297; 313; 334; 529; 777; 811; 934
--- rostratum: 445
--- sorokinianum: 448
--- turcicum: 219; 313
--- vagans: 573
--- victoriae: 3
Hemlock (see Tsuga)
Heterodera sp.: 122
--- glycines: 169
--- vigni: 345
Heteropogon contortus: a common wild grass in India is a host of Sclerospora sorghi and helps perpetuate the fungus which is a threat to maize production, 285
Hevea brasiliensis (rubber tree): Fomes root rot, retention of organic material on land prepared for replanting rubber will improve soil conditions, but should be confined to areas where root disease incidence is low, 295
Hibiscus esculentus (see Okra)
--- rosa-sinensis: a new ringspot virus disease was found in Nigeria which is different from any of the previously reported viruses on Hibiscus, 1040
High humidity chamber: a high humidity controlled temperature chamber will maintain humidity at 100% for obtaining consistent and reproducible infections by different pathogens, 814
Hobcaw Barony: a survey was made for the nematodes present in the 17,000 acres of this Barony located near Georgetown, South Carolina, 67
Hoplolaimus columbus: 369; 541; 641; 1000
--- galeatus: 822
Horned cucumber (Cucumis metuliferus): a potential source for resistance to watermelon mosaic virus 1 and squash mosaic virus, 735
Host resistance: four concepts for variety development and disease control are evaluated, 579
Hypoxylon atropunctatum: 622

Idaho: 75
Identification: of Cannabis cannabina and C. cannabis, 165
Illinois: 50; 411; 420
Indexing: of strawberry indicator clones for detection of 12 virus diseases, 28; of citrus trees with young tree decline and sand hill decline, 35; for grapevine fanleaf virus, 549; indexing lettuce seeds for seedborne lettuce mosaic virus by airstream separation of light from heavy seeds, 1037
India: 285; 345; 358; 767; 968; 1134
Indiana: 130; 430
Indicators: bark beetle attacks and discolored foliage were the best indicators of root rot in grand fir, 275
Infection sites: of Fusarium solani on yellow poplar seedlings, infections occurred through wounds on stems or through roots whether wounded or not, 754
Inhibition: of colonization by Ceratocystis ulmi in benomyl-treated elms, 511; of mycelial growth of several organisms by Trichoderma viride, 915
Inoculation: of germinating apple seedlings for testing pathogenicity of Erwinia amylovora isolates, 181; sawtooth oak seedlings inoculated with aeciospores

of Cronartium fusiforme proved to be hosts of the fungus, 198; of coniferous seedlings with Arceuthobium, on a continuing basis, 235; of apple rootstock selections with Erwinia amylovora, 243; zoospore inoculation of citrus stems with Phytophthora parasitica for determining relative susceptibility of citrus rootstocks to foot rot, 713; oak wilt development in red oaks after bole- or root inoculation, 738; of corn and sorghum with conidia of downy mildew fungus by method that is quantitative and more predictable than methods using conidia on leaf pieces, 825
Inoculum: soil assay developed to evaluate gladiolus cultivars for their tolerance to soilborne Fusarium in field samples, 184
Insect pests: of sweetpotato, carbofuran gave good control of sweetpotato flea beetle and southern potato wireworm, 239
Iowa: 448
Ipomoea setosa: grafted tissue from "hardcore" sweetpotato to Ipomoea setosa induced viruslike symptoms, 17
Iraq: 996
Irrigation water: occurrence of Phytophthora species and other potential plant pathogens in recycled irrigation water, 945
Isariopsis griseola: 330
Isolates: of barley yellow dwarf virus, mixed infections doubly infecting winter wheat and barley were common in New York in 6-year study, 472; no benomyl-tolerant isolates of Monilinia fructicola and M. laxa were found in California stone fruit orchards, 663
Israel: 1142
Italy: 691

Jamaica: 214
Japan: 603
Jelly melon (Cucumis metuliferus): two accessions were highly resistant to watermelon mosaic virus 1 and hypersensitive resistant to squash mosaic virus, 735
Johnsongrass (see Grasses)
Juniperus spp.: Sclerophoma pythiophila is reported for the first time on several Juniper species in Wisconsin and caused a tip dieback, 653
--- chinensis: benomyl gave best control of several fungicides for control of Phomopsis blight, 1012
--- virginiana (eastern redcedar): proximal spread of Fomes annosus in inoculated redcedar was significantly greater than in loblolly pine, 282; benomyl and thiophanate-methyl controlled Phomopsis blight on 2-year-old nursery seedlings, 476; Aureolaria grandiflora var. serrata, "false foxglove," was growing as parasite on J. virginiana in Lamar County, Texas, 773

Kansas: 267; 469; 566
Kentucky: 90; 188
Kenya: 1020

Leaf blister blight of tea (see Blight)
Leaf blotch: of oats, caused by Helminthosporium avenae, caused severe damage in Texas in 1973, 80
Leaf freckles: of corn, caused by Corynebacterium nebraskense, 113 corn lines were artificially inoculated to test for susceptibility, 956
Leaf graft: of strawberry, a modified technique by removal of all aerial organs except the grafted leaves improves detection of graft-transmissible diseases, 203
Leafhopper (Empoasca): suggested as the most likely insect vector of witches' broom of pigeonpeas, 96
Leaf necrosis: of spring and winter wheats in South Dakota, by Pseudomonas syringae, 1061
Leaf scald: of sugarcane, isolation of Xanthomonas albilineans from sugarcane leaves, 439
Leaf scorch: of almond, a newly described disorder of almond trees in California, 99
Leaf spot: zonate leaf spot of sorghum, caused by Gloeocercospora sorghi, in the northern sorghum belt, 267; of water hyacinth, caused by Cercospora piaropi, offers possibility of utilizing C. piaropi as a "biological herbicide," 277; of cherry, caused by Coccomyces hiemalis, was nearly completely controlled in greenhouse experiments with CW-524 (triforine), 326; angular leaf spot of bean, caused by Isariopsis griseola, discovered in Wisconsin 19 years after first appearance, 330; zonate leaf spot, caused by Cristulariella pyramidalis, was found on lima bean, snap bean and 'Concord' grape leaves, 421; of Aechmea fasciata, caused by Helminthosporium rostratum, 445; of Poa pratensis, caused by Helminthosporium sorokinianum, 448; gray leaf spot of corn, caused by Cercospora zeae-maydis in Virginia, 456; Cercospora zeae-maydis was the most destructive corn disease in the mountains of North Carolina, 532; target leaf spot of cucumber, caused by Corynespora cassiicola, controlled with benomyl, 636; of peanut, caused by Phomopsis sp., and was previously

unrecorded, 640; Cercospora leaf spot of peanut, disease forecasting facilitates chemical control, 666; of Rieger begonia, Xanthomonas begoniae stored for a year in dried leaves was still infective, 814; Cercospora leaf spot of sugarbeet controlled with fungicides, 952
---: of corn: reactions of 76 inbred lines to Helminthosporium leaf spot caused by a Helminthosporium not yet determined by name, 399; caused by a Helminthosporium isolate 72: 44-6, field reactions of 111 inbred lines to this isolate are listed, 909; 87 corn hybrids in the seedling stage varied in reaction to isolate 72: 44-6 (ATCC 24962), 975
Leaf-streak: of daylily, caused by Collecephalus hemerocalli, found in Pennsylvania, 102
Leaf wetness: a recorder was designed that records simultaneously temperature and leaf wetness on the same drum for immediate evaluation of all recorded wetness periods, to determine infection periods for apple scab, 133
Lebanon: 133
Lens culinaris: new experimental host of tobacco streak virus, 354
Lettuce: downy mildew, caused by Bremia lactucae, found in California head lettuce, 118; bottom rot controlled in New York tests by growing lettuce on ridges and spraying with benomyl or chlorothalonil, 837; separation of mosaic virus-containing seeds, 1037; lettuce mosaic accounted for over 90% of virus infections in commercial lettuce in Florida, 1087
Leucostoma persoonii: 971
Ligustrum japonicum: Cercospora leaf spot controlled with several fungicides, 1015
Liriodendron tulipifera (see Yellow poplar)
Litchi: a description of nine fungal postharvest diseases caused by Aspergillus spp., Cylindrocarpon tonkinense, Botryodiplodia theobromae, Colletotrichum gloeosporioides, and Pestalotia, 1134
Longevity: Xanthomonas begoniae, the cause of bacterial leaf spot of Rieger begonia, will survive for at least a year in dried leaves, 814
Longidorus elongatus: 591
--- macrosoma: 913
Lophodermium pinastri: 853
Losses: caused by potato virus Y infection in peppers, 141; in cucurbits in Morocco from watermelon mosaic virus-2, 143; in Florida cucumbers and peppers on retail market from decay, mechanical injury and chilling, 558
Lousiana: 201
Lycopersicon esculentum (see Tomato)

Macrophomina phaseolina: 50; 169; 1034
Macrosiphum avenae: 472
Maesopsis eminii (African hardwood tree): Fusarium solani and Volutella sp. are important canker-forming pathogens, 463
Maize (see Corn)
Male-sterile cytoplasm (see Cytoplasm)
Mango: powdery mildew, symptoms and morphology of the pathogen, Oidium mangiferae, 45
Maple: sugar, occurrence of an annual canker on sapling and pole-sized sugar maple trees in Minnesota, 1042
Market diseases: Sclerotinia rot, the most serious market disease of snap bean pods was controlled with chemical dips, 59; bacterial soft rot of pepper and mechanical injury caused the most loss in Florida bell peppers in Greater New York retail outlets, 367; of cucumber and pepper on the New York market, 558; the causal agent of a disease of sweet corn that causes a red discoloration of affected kernels was determined to be Epicoccum nigrum, 1065
Maryland: 727
Massachusetts: 483
Mating type: frequencies of mating type A in race T isolates of Bipolaris maydis in North Carolina, 529; mating types of oak wilt fungus, Ceratocystis fagacearum, in Arkansas, 622
Melampsora larici-populina: 423
--- lini: 311
--- medusae: 423; 721
Melanose: of grapefruit, was controlled on Texas grapefruit with several fungicides applied postbloom and 1 month later, 504
Melanospora sp.: 180; 373; 822
Meloidogyne spp.: 169; 369; 991; 1020
--- arenaria: 1075
--- hapla: 128; 594; 860
--- incognita: 88; 239; 361; 374; 517; 541; 562; 746; 749; 1048
"Melting out": Helminthosporium vagans was controlled on 'Merion' Kentucky bluegrass with single fall application of benomyl and PCNB, 573
Methods and techniques: three procedures were used for testing antagonism between ungerminated spores of Trichoderma viride and their effect on growth of Stereum purpureum, 71; simple method for purifying some plant viruses in laboratories without major facilities, 126; increment hammer and paste method of applying Trichoderma viride to trunk of peach trees was effective in decreasing silver leaf symptoms, 147; for production of healthy and diseased sugarbeet seedlings for host-pathogen investigations of Aphanomyces black root, 157; modified leaf grafting

technique by removal of all aerial organs but grafted leaves resulted in increased transmission, reduced incubation periods, and increased severity of initial symptoms of graft-transmissible diseases of strawberry, 203; method described for routinely producing coniferous seedlings infected with dwarf mistletoe on a year-round basis, 235; 10 Indian and American barley cultivars and an isolate of Helminthosporium gramineum were selected for evaluating intensity of infection against this pathogen, 265; simplified method for inducing asexual sporulation in Glomerella cingulata, 300; bark increment hammer was used to sample green stalks in corn stalk rot survey in Minnesota, 363; a method for assessing preharvest fungicide treatments on postharvest fruit rot of red raspberry, 391; isolation of Xanthomonas albilineans from sugarcane leaves, 439; forecasting method facilitates chemical control of Cercospora leafspot of peanuts, 666; measurement of water uptake as a diagnostic tool for diagnosis of young tree decline, blight and sand hill decline in citrus, 801; centrifugation-flotation method extracted more nematodes of most species from turf samples than the sugar-flotation-sieving method, 822; method for production of apothecia of Monilinia fructicola from brown rot-infected stone fruits for teaching purposes, 844; purification of cultures of wood decay fungi with benomyl-malt agar, 902; a photocopy system was developed to evaluate fluoride injury on gladiolus indicator plants in Ontario, 1091

---: assay: a rapid, sensitive biochemical assay for "Texas" cytoplasm in corn is described by use of Helminthosporium maydis, race T pathotoxin, 777; an assay method that detects Verticillium in safflower seed, 926; a bioassay technique was used to compare effectiveness of 17 organic and 3 copper fungicides in their ability to deposit on the tea leaf, ability to resist erosion and the stability of their biological activity, 928

---: equipment: a combined temperature-leaf wetness recorder for improved apple scab control that could be useful for other fungus control programs, 133; collection technique for ascospores of Whetzelinia sclerotiorum, 190; field evaluation of air pollution with Mandl chamber, 683; a high humidity and controlled temperature chamber of low cost that will maintain humidity at 100%, 814; indexing lettuce seeds for seedborne lettuce mosaic virus was aided by vertical airstream separation of light and heavy seeds, 1037

---: inoculation: a comparison of four methods for inoculating seeds of susceptible barley cultivars with a virulent strain of Helminthosporium gramineum, 32; needle inoculation of germinating open-pollinated apple seedlings is reliable for testing suspected Erwinia amylovora isolates, 181; spray inoculation technique compared favorably with oak-leaf technique for inoculating slash pine seedlings with fusiform rust, 584; zoospore inoculation of citrus stems with Phytophthora parasitica, 713; a quantitative technique for inoculating corn and sorghum with conidia of Sclerospora sorghi, 825

Metrosideros collina ssp. polymorpha (ohia): decline in Hawaii of the ohia tree and possible role of Phytophthora cinnamomi, 1069

Michigan: 180

Microsphaera sp.: on soybean cultivars in Georgia, 723

Milkweed vine (Morrenia odorata): Phytophthora citrophthora was isolated from roots and stems of dying milkweed vines in citrus groves in Florida; this isolate did not infect citrus, suggesting race differences in P. citrophthora, 355

Minnesota: 363; 693; 1042

Mississippi: 165

Missouri: 92

Mistletoe: dwarf: coniferous seedlings were successfully inoculated with Arceuthobium seeds which had been stored, 235; was abundant in mixed conifer stands at lower elevations on the Fort Apache Indian Reservation, Arizona, 418; removal of all bole-infected trees below 5 inches in diameter at bole infection center will eliminate nearly all bole infections capable of spreading infection, 1066

Momordica charantia: a common weed in citrus orchards is a host of Radopholus similis, 561

Monetary loss: quality and quantity reductions in mushrooms from three fungal diseases, Verticillium malthousei, Mycogone perniciosa and Dactylium dendroides, in commercial mushroom houses in Pennsylvania, were valued at $9.1 million, 987

Monilia fructigena: 915

Monilinia fructicola (brown rot of peach): 208; 663; 844

--- laxa: 663; 915

--- vaccinii-corymbosi: 840

Morocco: 141; 143; 252; 307; 309; 1117; 1121; 1123

Morphology: of Oidium mangiferae, causal

organism for powdery mildew of mango, 45; of Cannabis cannabina and C. cannabis, 165; of Phytophthora megasperma var. sojae, 525
Mowing height: Kentucky bluegrass cultivars vary in their reaction to mowing height and fertilization in relation to infection from Fusarium blight and Sclerotinia dollar spot, 514
Mummy berry: of highbush blueberry, triforine controlled Monilinia vaccinii-corymbosi primary infection and increased yields over control, 840
Mushrooms (Agaricus bisporus): losses from three mushroom diseases in commerical mushroom houses in Pennsylvania amounted to over 9 million dollars, 987
Muskmelon (see Cucurbits)
Mycogone perniciosa: 987
Mycological description: of Phialophora fastigiata, the cause of vascular wilt of Michaelmas daisy, 57; of Sydowia polyspora, the cause of tip dieback on white fir, 94; of Melampsora larici-populina and M. medusae, the cause of poplar rust, 423
Mycoplasma: witches' broom disease of pigeonpeas in Puerto Rico, is caused by mycoplasmalike organisms, 96; evidence indicates that mycoplasmalike organisms were associated with tomato big bud, 211; media supporting Spiroplasma citri did not support the growth of the mycoplasmalike agents causing aster yellows and corn stunt diseases, 797; sieve elements of Canada thistle showing typical yellows symptoms contained mycoplasmalike bodies, 906; detection of motile spiroplasma filaments in abdominal smears and in hemolymph removed from corn stunt-infected individuals of the insect vector Dalbulus maidis, 1109; mycoplasmalike bodies found in tulip in Israel, 1142
Mycosphaerella citrullina: 558; 899
Myzus persicae: 141; 1123

Naemacyclus niveus: 287
Nebraska: 267; 782
Nectria cinnabarina: 872
--- coccinea var. faginata: 718
Needlecast: of Scots pine, in Pennsylvania, of epidemic proportion, 287; Lophodermium pinastri in Scotch pine Christmas trees was controlled with maneb, a mixture of chlorothalonil and cycloheximide, and chlorothalonil, 853; of blue spruce Christmas trees, caused by Rhizosphaera kalkhoffii, 1094
Nematicides: several tested together: for control of nematode and insect damage and on yield and grade of sweetpotato, 239; none tested was effective for control of Cylindrocladium black rot of peanuts, 348; for control of root-knot and reniform nematodes in potato, 517; for control of Columbia (lance) nematode on soybeans, 641; phenamiphos and ethoprop reduced root galls of Meloidogyne incognita more than other chemicals tested, 749; for control of Meloidogyne hapla in peony, 860; 997
---: aldicarb: for control of reniform nematode on cotton, 337; was more effective than other systemic nematicides tested for control of root-knot and reniform nematodes in potato, 517; effectively controlled root-lesion nematodes in greenhouse roses and gave some control of the two-spotted mite, Tetranychus urticae, 730
---: BAY 68138 (see phenamiphos)
---: carbofuran: was best of chemicals tested for control of sweetpotato flea beetle and southern potato wireworm, 239; gave good results for control of lance nematodes on soybeans, 641; increased grade and yield of cantaloup and controlled Meloidogyne incognita, 746; controlled M. incognita, on lima bean, 749
---: 1,3-D: 1,3-D and DBCP fumigation reduced root populations of Hoplolaimus columbus and Meloidogyne incognita in cotton, 541; applied at high doses and with deep placement and split-depth application to replant vineyard soils, 1,3-D gave promising nematode control, 562; 1,3-D in combination with ultra-high frequency electromagnetic energy (UHF) controlled reniform nematodes and common purslane in cowpea field, 985
---: DBCP: lint yield and stand count of cotton were highest after fumigation with DBCP; herbicides (prometryn and trifluralin) had no apparent nematode control, 120; reduced root population of Hoplolaimus columbus and Meloidogyne incognita in cotton, 541; effective in control of H. columbus in South Carolina, 641; gave control of M. hapla in peony but failed to kill all nematodes in fleshy storage roots, 997
---: DD: controlled root-knot nematodes in sweetpotato, 239; tested for control of Rotylenchulus reniformis on cotton, 337; reduced nematode populations and reduced levels of Verticillium albo-atrum in potato fields, 374; controlled Columbia (lance) nematode on soybeans in S. C., 641; controlled Meloidogyne incognita and increased cantaloup grade and yield, 746;

Nematicides (continued): apple rootstocks, differed in growth response to soil fumigation, 1007; DD and EDB reduced the numbers of Meloidogyne incognita and M. javanica larvae in bean plots, 1020; fumigation reduced nematode populations and resulted in increased yield of sweetpotato in Trinidad, 1048
---: EDB; yield and grade of sweetpotatoes was increased by soil fumigation for control of root-knot nematodes, 239; yield and grade of cantaloup were significantly increased and Meloidogyne incognita was controlled with EDB soil treatment, 746; M. incognita and M. javanica were reduced in number in bean plots by EDB and DD, 749
---: ethoprop: ethoprop and carbofuran controlled Meloidogyne incognita and more jumbo and No. 27 grades of cantaloup were produced than by other nematicide treatments, 746; ethoprop and phenamiphos applications controlled M. incognita and increased yields of lima bean, 749
---: fensulfothion: increased yield and grade of cantaloup and controlled M. incognita, 746; reduced number of root galls of M. incognita on lima bean and increased yields, 749
---: MBR, deep and shallow placements of soil fumigants were tested to control reniform nematode on cotton in Texas, 337
---: oxamyl: Longidorus elongatus was controlled on mint with oxamyl foliar spray, 591; at 30°C, oxamyl was effective against Pratylenchus penetrans, 708; gave only fair control of Pratylenchus penetrans on greenhouse roses, 730; sharply reduced citrus nematode populations by foliar application, 882; as a monthly spray at 1/4 lb a.i./gal, oxamyl was an effective control for Aphelenchoides fragariae on Rieger begonia, 911
---: phenamiphos (BAY 68138): controlled root-knot nematode on sweetpotato, 239; gave good control of lance nematodes on soybeans, 641; practically eliminated Pratylenchus penetrans in untreated, unplanted pots, 708; controlled Meloidogyne incognita and increased yield and grade of cantaloup, 746; gave good control of M. incognita and increased yield of lima beans, 749; controlled M. arenaria on gardenia and fig plants, 1075
---: SD-7727, gave high degree of control for root-knot nematode in peony, 997
---: thionazin, as a dip on bare roots controlled Meloidogyne hapla, 997
Nematodes: a Verticillium wilt-nematode complex hypothesized as possible cause of early dying disease of potatoes in north Florida, 374; more nematodes of most species were extracted from turf samples by the centrifugation-flotation method than by the sugar-flotation-sieving method, 822; control of reniform nematode in cowpeas by soil application of ultra-high frequency electromagnetic energy, 985
---: burrowing, Momordica charantia is a new host of Radopholus similis, 561
---: citrus nematode, foliar applications of oxamyl suppressed citrus nematode in Texas orchards, 882
---: cyst, Heterodera cajani, caused malformations in cowpea, 1130
---: dagger: 1,3-D at high dosage and deep placement controlled Xiphinema americanum in replant vineyards, 562; controlled in California prune orchards, 633
---: foliar: Aphelenchoides fragariae were controlled on Rieger begonia with oxamyl, 911
---: lance: Hoplolaimus columbus, survey of 306 cotton and soybean fields in east Georgia revealed low populations of lance nematodes, 369; DD and DBCP provided better control than other nematicides for control of the Columbia (lance) nematode on soybeans, 641; host range of Hoplolaimus columbus, 1000
---: needle: Longidorus elongatus was controlled on peppermint with oxamyl, 591; DBCP drenches to glasshouse roses killed L. macrosoma, 913
---: pin, Paratylenchus sp. were controlled in replant vineyards with 1,3-D at high dosage and deep placement, 562
---: plant-parasitic: 16 genera of stylet-bearing nematodes were found in a survey of Hobcaw Barony, 67; Pratylenchus zeae, Trichodorus minor, Tylenchorhynchus martini, and Heterodera sp. are reported for first time from Trinidad, 122; plant-parasitic nematodes, pathogenic fungi, and endomycorrhizal fungi associated with soybean roots in Florida, 169; 1,3-D at high dosage and deep placement gave control in grape replant vineyards in California, 562; nematode populations in California prune orchards were comprised of pin, dagger, ring and root-lesion nematodes, 633; the previous crop as well as the geographic area in N. C. greatly affected nematode occurrence and activity, 991; in Trinidad, soil fumigation with DD resulted in increased sweetpotato yields, 1048

---: reniform: control of Rotylenchulus reniformis and Meloidogyne incognita on potatoes with systemic nematicides, 517; of R. reniformis with combination soil treatment of ulta-high frequency electromagnetic energy and 1,3-D in cowpea field, 985
---: ring: Criconemoides xenoplax, its possible role in short life of peach, 76; in California prune orchards, 633
---: root-knot: overwintering of Meloidogyne incognita in root galls of sugarbeet in California, 88; M. incognita acrita, interaction of nematicides and herbicides for pest control on cotton, 120; two Pacific Northwest biotypes of M. hapla produced on corn and oat, 128; inheritance of resistance to M. incognita in cowpea, 361; M. incognita was controlled in cantaloup plots and yields of melons were increased, 746; controlled on lima beans, 749; M. hapla controlled in peony, 860; in peony by dips with thionazin, 997; M. incognita and M. javanica were controlled with DD and EDB in bean fields in Kenya, 1020; M. arenaria controlled on gardenia and fig plants with phenamiphos, 1075
---: root-lesion: comparative tolerance of walnut species, walnut hybrids, and wingnut to Pratylenchus vulnus, 630; in California prune orchards, 633; treatment of Pratylenchus penetrans with aldicarb and oxamyl on roses, 730; P. penetrans caused root damage to apple trees in New York, 1007
---: stubby-root, Trichodorus allius and T. teres are vectors of tobacco rattle virus affecting potato production in Oregon, 269
New Brunswick: 872
New diseases: leaf scorch of almond, also referred to as "almond decline" and "golden death," 99; new leaf disease of Aechmea fasciata is caused by Helminthosporium rostratum, 445; of asparagus in New Zealand, caused by Phytophthora megasperma var. sojae, 525; a new viruslike disease found on tomato in eastern Washington, 965
New distribution: tomato spotted wilt virus found on peanut in Texas, 23; pear blast in Connecticut, 40; Phialophora fastigiata found in New Zealand as a vascular wilt pathogen of aster, 57; of oak wilt in Missouri, 92; of tip dieback fungus, Sydowia polyspora on white fir in California, 94; of leaf-streak of daylily, in Pennsylvania caused by Collecephalus hemerocalli, 102; first record for Cercospora cannabina on Cannabis sativa in North America in Mississippi, 1972, 135; several fungi reported for first time on highbush blueberry in Michigan, 180; of Alternaria macrospora as a pathogen on cotton in Louisiana, 201; big bud of tomato reported from the Eastern U.S., 211; of zonate leaf spot on sorghum, in Nebraska and Kansas, 267; of Cercospora piaropi, that causes a leaf spot on water hyacinth, 277; first record of Ascochyta rabiei in North America, in Saskatoon, Saskatchewan, Canada, 342; of phloem necrosis in New Jersey in 1973, 387; maize chlorotic dwarf virus found in Illinois on johnsongrass, 420; black shank of tobacco and Pseudomonas syringae on blighted pear blossoms were found in Connecticut, 428; of sorghum downy mildew in Indiana, 430; potato pink root reported for the first time from Delaware, 443; brown root rot of tomato found in Massachusetts, 483; Helminthosporium hawaiiense is reported for the first time from the continental United States on Rhodesgrass in Florida, 490; Endothia parasitica found on live oak in North Carolina and Florida, 596; Cylindrocladium floridanum in New Zealand, 705; cotton wilt in Iraq, 996; of needlecast disease of blue spruce, caused by Rhizosphaera kalkhoffii, found in Minnesota, Wisconsin, Michigan and Indiana, 1094
New hosts: of Armillaria mellea in Hawaii, 102; several fungi are reported for the first time on highbush blueberry, 180; Cronartium fusiforme, sawtooth oak found susceptible, 198; lima bean, snap bean and Vitis labrusca are new hosts of Cristulariella pyramidalis, 421; oat is new host of Drechslera siccans in Argentina, 507; pepper is a new host of Fusarium equiseti and Myrothecium verrucaria in Argentina, 507; bean is new host of Fusarium equiseti in Argentina, 507; Fraxinus americana, infected by tobacco mosaic virus, is first evidence of TMV actually causing a disease of a tree, 536; Momordica charantia is host of burrowing nematode that causes a serious disease of citrus, 561; Sclerophoma pythiophila found in Wisconsin on Juniperus species and is cause of a tip dieback, 653; Galinsoga parviflora is susceptible to Whetzelinia sclerotiorum and could play an important role in the survival and distribution of the pathogen, 700; Aureolaria grandiflora var. serrata, was found growing on Juniperus virginiana, 773; sunflower

is a new host for Xanthomonas sp., Phoma herbarum, and Cochliobolus sp. in Tanzania, 896; red fescue was found infected with Puccinia pygmaea, 1028
New Jersey: 387; 683
New South Wales: 437; 845
New York: 211; 367; 784; 837; 1007
New Zealand: 57; 423; 525; 705
Nigeria: 616; 1080
North Carolina: 282; 318; 348; 439; 521; 529; 532; 596; 822; 991

Oak: Oak sawtooth (Quercus acutissima), is a new host of Cronartium fusiforme, 198; growth loss of scarlet oak (Quercus coccinea) due to oak decline in Virginia, 396; live oak (Quercus virginiana) in North Carolina and Florida were found infected with chestnut blight, 596; California black oak (Quercus kelloggii) proved to be susceptible to southern fusiform rust, 497; Quercus nigra, oak leaf method for inoculating pine seedlings with fusiform rust was compared with spray inoculation method, 584
---: oak wilt: protection of sand-grown red oak seedlings from oak wilt disease by drenching with benomyl, 65; first report of oak wilt fungus being isolated from two counties in Missouri, 92; in red oaks that were root-inoculated with Ceratocystis fagacearum, 738
Oat: sensitivity of leaf tissue to wilt toxin victorin, and calcium content, 3; oat leaf blotch caused severe damage to oats in central Texas, 80; Meloidogyne hapla reproduced on 'Lee' oats in the Pacific Northwest, 128; early teliospore formation is a trait of oat line 66B1285 when attacked by Puccinia coronata avenae, 467; new record of Fusarium moniliforme on oat seed in Argentina, 507; races of Puccinia graminis f. sp. avenae present in the United States in 1973 are summarized, 605; susceptibility of leaves to Cylindrocladium, 1017
Ohia (Metrosideros collina ssp. polymorpha): distribution of Phytophthora cinnamomi and its possible role in the decline of ohia trees in Hawaii, 1069
Ohio: 320
Oidium sp.: 12
--- begoniae (imperfect stage of Erysiphe): 875
--- mangiferae: 45
Oklahoma: 687
Okra (Abelmoschus esculentus = Hibiscus esculentus): virus-induced mosaic disease of okra in Nigeria, 616
Onion: white rot, caused by Sclerotium cepivorum, controlled by several fungicides by furrow and root-dip applications, 6; high incidence of onion yellow dwarf virus in Morocco, 252
Ontario: 677; 1091; 1105
Oregon: 75; 766
Overwintering: of Meloidogyne incognita in root galls of sugarbeet in California, 88
Ozone: ten bean varieties rated for relative susceptibility to ozone, 14; ozone injury was reduced on Poa annua by benomyl and thiophanate-methyl when applied as a soil amendment or as a soil drench, 41; 1,4-oxathiin derivatives protect plants against ozone, 162; injury to current needles of ponderosa pine was greatest when needles were still elongating, 660

Pacific Northwest: 128; 492; 766
Papulospora sp.: 180
Paratylenchus neoamblycephalus: 633
Pea: reappearance of common wilt of pea in Washington, caused by Fusarium oxysporum f. sp. pisi race 1, after almost 40 years, 62; epiphytotics of pea anthracnose in 1972 and of downy mildew in 1973 were reported from Wisconsin, 226; behavior of Fusarium solani f. sp. pisi and F. solani f. sp. phaseoli was tested individually and in combination on peas and beans, 500; nematode and purslane control by UHF electromagnetic energy and soil fumigation, 985
Peach: short life of peach trees in Georgia appears to be closely associated with site, nematodes, bacterial canker, and cold injury, 76; curative treatment of peach trees against silver leaf disease (Stereum purpureum) with Trichoderma viride, 147; peach brown rot and scab, were most effectively controlled with benomyl and BAS 3201 F, 208; rusty spot, HOE 2873 was highly effective for control, 254; peach tree decline, postpruning sprays not effective in control, 388; stem pitting discovered in Georgia in 1973, 553; Leucostoma canker, development of cankers and distal foliar symptoms are not influenced by frequency of irrigation, 971
Peanut (Arachis hypogaea): tomato spotted wilt virus was found affecting peanuts in south Texas and is a first report of TSWV disease of peanut in the U. S., 23; peanut mottle virus was found in each of 117 peanut fields surveyed in Georgia, 107; seedling and pod rot disease, caused mainly by Rhizoctonia solani, was controlled in Egypt with

new systemic fungicide TCMTB-60W, 176; 138 new cases of Cylindrocladium black rot were reported from North Carolina during 1973, 348; a previously unrecorded leaf spot of peanut, caused by Phomopsis sp., was found in India, 640
Pear: pear blast: caused by Pseudomonas syringae was reported for the first time from Connecticut, 40; was identified in Chile, 568
Pecan: benomyl and thiophanate-methyl controlled pecan scab (Fusicladium effusum), 687; nut-rotting fungus Botryosphaeria ribis was isolated from eight pecan cultivars, 1030
Pectobacterium carotovorum: 214
Penicillium digitatum: 933; 1143
Pennsylvania: 102; 287; 718; 719; 934; 987
Peony: fumigated soil followed by annual sidedressings of systemic or contact nematicides effectively controlled Meloidogyne hapla in peony, 860; bare-root dip with thionazin controlled M. hapla, 997
Pepper (Capsicum annuum): serious losses in peppers in Morocco from potato virus Y, 141; bacterial soft rot and mechanical injury are the cause of most of the culls of Florida bell peppers at New York City Terminal Market, 367; Drechslera sorokiniana and Fusarium moniliforme found on pepper seeds in Argentina, 507; market diseases on the New York market were caused chiefly by Alternaria rot and gray mold, mechanical injury and chilling, 558; potato virus Y and tobacco etch virus singly or together were the most frequent viruses isolated from commercial peppers in California, 1002
Peppermint (Mentha piperita): oxamyl controlled Longidorus elongatus in greenhouse experiments, 591
Peregrinus maidis: 122
Peronospora pisi: 226
Pestalotia sp.: 1134
Petunia: growth alterations associated with air pollution, 683; dodder was found for the first time in Bermuda, on petunia, 974
pH: resistance or susceptibility of tomato and alfalfa plants to Meloidogyne hapla was not appreciably affected by pH level, 594; preliminary data suggest that soil in the vicinity of western hemlock infected with Fomes annosus is lower than that of soil around healthy trees, 1077
Phaseolus lathyroides: susceptibility of P. lathyroides, a potential forage plant in Brazil, to soybean mosaic virus could make it a permanent source of the virus, 322
--- vulgaris (see also Bean): 125 processing bean cultivars and breeding lines were tested for reaction to bean root rot complex in Wisconsin, 229; several dry edible bean varieties were found to exhibit tolerant reaction to white mold epidemic in western Nebraska, 782
Phialophora fastigiata: 57
Philippines: 544
Phloem necrosis: of elm, recent discovery in New Jersey, 387
Phomopsis sp.: 173; 180; 414; 416; 640; 978
--- juniperovora: 476; 1012
--- viticola: 915
Phymatotrichum omnivorum: 373
Physalis floridana: is a good production host for Satsuma dwarf virus, 603
Phytophthora spp.: 945
--- cinnamomi: 318; 437; 650; 1069
--- citrophthora: 355; 945
--- drechsleri: 703
--- erythroseptica: 443
--- megasperma var. sojae: 221; 352; 525; 789
--- nicotianae var. nicotianae: 767
--- palmivora: 1100
--- parasitica: 669; 713; 945
Picea engelmannii (Engelmann spruce): was parasitized by dwarf mistletoe at elevations below 10,000 ft on the Fort Apache Indian Reservation, Arizona, 418
--- pungens (blue spruce): Rhizosphaera kalkhoffii, the cause of needlecast was found on blue spruce in North Carolina, 439; found on the blue spruce Christmas tree plantations; 1094
Pierce's disease: electron microscopic evidence of Pierce's disease in six grape vines at College Station, Texas revealed presence of rickettsia-like bacteria, 780
Pigeonpea (Cajanus cajan): witches' broom disease, caused by a mycoplasmalike organism, is present in Puerto Rico, 96
Pink rot: of potato, caused by Phytophthora erythroseptica, was reported for the first time from Delaware, 443
Pinus spp.: Cylindrocladium floridanum has been isolated from rotting and dying seedlings of pine in New Zealand, 705
--- banksiana (jack pine): first record of Discosia on a coniferous host, 677
--- contorta (lodgepole pine): removal of trees below 5 inches having dwarf mistletoe bole infections will eliminate nearly all infections capable of spreading dwarf mistletoe, 1066
--- elliottii var. elliottii (slash pine): incidence and impact of pitch canker in slash pine plantations was measured

for 3-year period, 451; basidiospore spray inoculation of pine seedlings with Cronartium fusiforme produced response similar to oak leaf inoculation method, 584; systematic evaluation was made of incidence of fusiform rust in slash and loblolly plantations in the South, 1137
--- jeffreyi (Jeffrey pine): is highly susceptible to fusiform rust, 497
--- nigra (Austrian pine): Pinus nigra seedlings found susceptible to Cronartium comandrae, 616
--- ponderosa (ponderosa pine): infection technique developed to infect pine seedlings with dwarf mistletoe on a year-round basis, 235; because of high susceptibility of ponderosa pine, P. jeffreyi, P. radiata, and alternate host, California black oak to fusiform rust, care must be taken that the rust is not introduced to the Western U. S. A., 497; current needles, still elongating, showed the greatest sensitivity to ozone, 658
--- radiata (Monterey pine): this western pine is highly susceptible to fusiform rust, 497
--- resinosa (red pine): Scleroderris canker was controlled with chlorothalonil and chlorothalonil + cycloheximide, 1097
--- strobus (eastern white pine): use of high nitrogen fertilizer lessened symptoms of air pollution on white pine Christmas trees in Virginia, 150
--- sylvestris (Scots or Scotch pine): Naemacyclus niveus is cause of needlecast epidemic in Pennsylvania, 287; control of Lophodermium needlecast was achieved with fungicides, 853; survival of Gremmeniella abietina (Scleroderris lagerbergii) in marketed Christmas trees, 892
--- taeda (loblolly pine): spread of Fomes annosus in inoculated loblolly pine was significantly less than in eastern redcedar, 282; greenhouse screening of loblolly pine of known parentage for resistance to Fomes annosus, 409; incidence of fusiform rust in Southern States was evaluated, 1137
Plant architecture: modification of plant habit or structure in 'Great Northern' dry beans affected the severity of white mold infection, 379
Planting site: for peach, its importance in the complex that causes short life of peach trees, 76
Plant quarantine: long smut (Tolyposporium ehrenbergii), a pathogen that does not occur in the United States was found on imported sorghum seed, 810
Poa annua (see Grasses)
--- pratensis (see also Grasses): occurrence of midsummer Helminthosporium sorokinianum leaf spot in Iowa is thought to be related to irrigation practices and possible changes in virulence in the pathogen, 448
Podagrica uniformis: 616
Podosphaera leucotricha: 625; 949
Poinsettia: air pollution effects on green weight, 683
Polymyxa graminis: 878
Poor drainage: suggested as a predisposing factor for weakening resistance of flowering dogwood to invasion by Armillaria mellea, 719
Poplar (see Populus)
Populus (poplar): New Zealand has recently experienced outbreaks of two rust species pathogenic to poplar, 423; breeding for resistance to Melampsora medusae, 721
--- tremuloides (aspen): a fresh blowdown in a Colorado aspen stand indicated a strong relationship between windthrow and butt-rotting fungus, Fomes applanatus, 110
Portulaca oleracea (common purslane): this weed and reniform nematodes were controlled in field planted to southern peas by a combination treatment with 1, 3-D and ultra-high frequency electromagnetic energy, 985
Postharvest decay: DCNA and thiabendazole controlled Sclerotinia rot of snap bean pods during storage, 59; causes of cullage of Florida bell peppers in New York wholesale and retail markets, 367; of red raspberry, assessment of preharvest fungicide treatments, 391; of litchi fruits in India, by nine different pathogens, 1134; of grapefruit inoculated with Penicillium digitatum, 1143
Potato: potato wart, caused by Synchytrium endobioticum, has been eradicated from West Virginia, 291; air pollution injury of potato plants grown in a growth chamber, 304; early dying disease in north Florida is possibly caused by a Verticillium wilt-nematode complex, 374; pink rot of potato, caused by Phytophthora erythroseptica, found in Delaware, 443; root-knot and reniform nematodes controlled with aldicarb, 517; effect of soil temperature on severity of Rhizoctonia solani infection on potato shoots, 646
Powdery mildew: of Antirrhinum, caused by Oidium sp., controlled with fungicides, 12; of mango, caused by Oidium mangiferae, 45; of apple, caused by Podosphaera leucotricha, 51 apple

cultivars rated for susceptibility throughout the United States and Canada, 625; of cucumber, chlorothalonil and zinc + maneb gave satisfactory control, 636; on Glycine max, caused by Microsphaera sp., 723; of Rieger begonia, susceptible cultivars were afforded protection by several fungicides, 875; of cantaloup, a high level of resistance to Sphareotheca fuliginea race 2 is present in Taiwan cantaloup introduction, 899; of sugarbeet, an epiphytotic occurred on sugarbeets in California in 1974, 904; Off-Shoot O and Off-Shoot T were effective as eradicants of mildew from dormant vegetative apple buds, 949
Pratylenchus sp.: 991
--- spp.: 169; 633
--- penetrans: 708; 730; 1007
--- vulnus: 630
--- zeae: 122
Proboscidea louisianica: new host of tobacco streak virus, 354
Prune: plant-parasitic nematodes in California prune orchards, 633
Pruning: winter pruning was more effective than fall pruning or postpruning sprays in control of peach tree decline, 388
Pruning date: for peach, early pruning made trees more susceptible to cold damage but was less important than site, presence of nematodes, bacterial canker, and cold injury, 76
Prunus (see also common names): no benomyl-tolerant isolates of brown rot fungus, Monilinia fructicola and M. laxa, were found in stone fruit orchards in California, 663
Pseudomonas sp.: 411
--- oryzicola: 1126
--- solanacearum: 86; 793
--- syringae: 40; 76; 388; 428; 568; 1061
Pseudoperonospora cubensis: 899
Pterocarya stenoptera (Chinese wingnut): was highly tolerant of Pratylenchus vulnus in California orchard trial, 630
Puccinia anthirrhini: 733
--- coronata avenae: 467
--- graminis: 459; 469
--- --- avenae: 467; 605
--- --- f. sp. tritici: 608; 806; 1085
--- polysora: 532
--- pygmaea: 1028
--- recondita: 272; 459; 469
Puerto Rico: 96
Pullularia spp.: 180
Purslane (see Portulaca)
Pyrenochaeta sp.: 180
--- lycopersici: 483
Pyricularia oryzae: 544
Pythium spp.: 169; 229; 373; 941
--- ultimum: 221; 774
Quarantine: potato wart quarantine in West Virginia has been lifted, 291
Quercus (see Oak)
Races: new race of flax rust discovered in North America, 311; race 2 of Fusarium oxysporum f. sp. lycopersici was recovered from greenhouse soil in Ohio to a depth of 90 cm, 320; maintenance of stock cultures of Helminthosporium maydis races T and O, 334; races 1 and 2 of Fusarium oxysporum f. sp. conglutinans and susceptibility of Crambe spp. and Brassica carinata, 479; the population of race T of Bipolaris maydis in North Carolina has declined suggesting that race T is unable to survive in N.C. on corn with normal cytoplasm, 529; of Helminthosporium maydis and H. carbonum on corn hybrids in normal cytoplasm in Pennsylvania in 1973, 934
---: Phytophthora megasperma var. sojae: race 4 is proposed, 352; physiologic races 3 and 4 were isolated from soybean for the first time in Indiana, 789
--- Puccinia: Puccinia graminis f. sp. avenae races in the United States in 1973, 605; Puccinia graminis f. sp. tritici races in the United States in 1973, 608; there is evidence of two populations of wheat stem rust in southern and northern areas of U.S. wheat growing areas, 806
Radopholus similis: 561
Raspberry: postharvest decay from Botrytis cinerea, 391; 144 cultivars and selections showed variations in susceptibility to spur blight, caused by Didymella applanata, 1024
"Red kernel": of sweet corn on retail market was caused by Epicoccum nigrum, 1065
Red stripe disease: the johnsongrass-infecting strain of sugarcane mosaic virus was identified as the cause of red stripe disease of sorghum in New South Wales and was found to be serologically related to virus causing maize mosaic, 832
Relative humidity: its effect on sporulation of Helminthosporium maydis on corn, 297
Remote sensing: previsual diagnosis of young tree decline of citrus by remote sensing infrared thermometer was not successful, 793
Resistance: inheritance pattern for resistance to root-knot nematode in cowpea, 361; resistance in tomato fruit to Rhizoctonia soil rot is thought to be

polygenic, 406; screening for resistance to Fomes annosus in loblolly pine, 409; Cercospora beticola resistance to benomyl fungicide, 434; resistance to Phytophthora root rot in Rhododendron species and hybrids, 650; in 'V 20' tobacco to tobacco etch virus, 658; of apple cultivars to cedar-apple rust and quince rust, 696; a white-seeded snap bean line 1273 was found to be highly resistant to seed decay and damping-off, 774; of Rieger begonias to powdery mildew, Schwabenland Red and related cultivars are susceptible, Aphrodite types and cultivars were all immune, 875; several wheat cultivars available in Florida have resistance to soilborne wheat mosaic virus, 878; to three cantaloup diseases in cantaloup PI 321005 from Taiwan, 899

---: to Dutch elm disease: of hybrids of Chinese elm and several elm species to Ceratocystis ulmi, 727; in New York study, 16 American elm trees from a collection of over 21,000 have resistance, 784

Rhamnus frangula ('Tallhedge'): freezing stress appears to predispose Tallhedge to attack by Tubercularia ulmea, that causes a wilting and dieback of stems, 937

Rhizina undulata: 492

Rhizobium japonicum: fewer root nodules were produced on soybean infected by Phytophthora megasperma var. sojae race 3 alone, or in combination with Pythium ultimum, 221

Rhizoctonia spp.: 114; 169

--- solani: 229; 373; 406; 636; 646; 941

Rhizopus spp.: 391

Rhizosphaera kalkhoffii: 439; 1094

Rhode Island: 848

Rhododendron: considerable variation exists among Rhodendron species in susceptibility to Phytophthora root rot, 650

Rhopalosiphum maidis: 1121

--- padi: 472

Rice: Cercospora oryzae and Trichoconis padwickii on rice seeds are new records for Argentina, 507; daily periodicity of conidia release of Pyricularia oryzae over a rice blast nursery was measured, 544; tungro virus, alternate hosts include Paspalum distichum, Echinochloa colonum, Cyperus rotundus and Cynodon dactylon, 856

Ridging: of lettuce plus sprays of benomyl or chlorothalonil was effective against infection by Rhizoctonia solani, 837

Root dips: control of Meloidogyne hapla on peony with bare-root dip in thionazin, 997

Rose: nematode control of greenhouse roses with aldicarb, 730; drench with DBCP killed Longidorus macrosoma on 'Dr Verhage' glasshouse roses, 913

Rot: bacterial rhizome rot of banana, 214

---: basal stem rot of coconut, caused by Ganoderma boninense, in Sri Lanka, 293

---: boll, losses of cotton by parasitic diseases in 1973, 373

---: Botryosphaeria ribis was isolated from eight pecan cultivars, 1030

---: bottom rot of lettuce, caused by Rhizoctonia solani, was controlled by ridged plantings and fungicide sprays, 837

---: brown root rot of tomato, first confirmed report of this disease on tomato in North America, in Massachusetts, typical brown root rot symptoms were reproduced on 22 tomato cultivars, 483

---: Cercosporella foot rot of wheat, influence of seeding date, varietal resistance, and benomyl spray on control, 554

---: corm, of gladiolus, assay of soilborne inoculum potential, 184

---: crown, of Gypsophila paniculata (baby's breath), caused by Phytophthora parasitica, 669

---: Cylindrocladium black rot of peanuts, incidence and control in North Carolina in 1973, 348

---: Fomes lignosus, infection of Hevea roots, 295

---: foot rot of citrus, caused by Phytophthora parasitica, zoospore-inoculation technique for determining susceptibility in citrus rootstocks, 713

---: fruit: of red raspberry, assessment of preharvest fungicide treatments on postharvest fruit rot by two techniques, 391; tomato, evaluation for resistance to Rhizoctonia soil rot, 406

---: peach brown rot: 208; no tolerant isolates of Monilinia fructicola were found in California stone fruit orchards, 663; production of apothecia, 844

---: Phytophthora root rot, caused by P. megasperma var. sojae, 95 soybean germplasm strains were resistant to races 1, 2, 3, and 4, 789

---: Rhizina root rot, in an old-growth conifer stand in the Pacific Northwest, 492

---: Rhizopus rot of peach, not controlled by any of six systemic fungicides tested, 208

---: root: several new hosts of Armillaria mellea have been found in Hawaii, 102; of quaking aspen, caused by Fomes applanatus, 110; of black walnut, caused by Cylindrocladium scoparium, 188; of soybean, effect of Rhizobium japonicum and Endogone mosseae on

soybean root rot, 221; root rot complex of processing beans, only four of 125 lines showed promising tolerance to the disease, 229; of grand fir (Abies grandis), 275; of corn seedlings as influenced by Helminthosporium leaf blights, 313; of Fraser fir Christmas trees in North Carolina, 318; of milkweed vine, caused by Phytophthora citrophthora, 355; Fomes annosus in giant sequoia, 478; of pea and bean, caused by Fusarium solani f. sp. pisi and F. solani f. sp. phaseoli, respectively, 500; of cassava, is caused by Phytophthora drechsleri, 703; of bean, was widespread throughout sampled area in southwestern Colorado, 890; of bean, BAY 22555 gave most promising results of fungicides tested, for control, 941

---: Sclerotinia rot of snap bean pods, controlled with DCNA and thiabendazole dips, 59

---: sheath-rot of rice, caused by Acrocylindrium oryzae, 358

---: shoot rot and leaf fall of cashew (Anacardium occidentale), was caused by Phytophthora nicotianae var. nicotianae, 767

---: stalk rot of corn, cob and shank inoculations with Diplodia maydis, 1113

---: white, of onion, controlled in Washington with fungicides, 6

Rotylenchus reniformis: 201; 259; 337; 369; 517; 985; 1048

Rubber (Hevea brasiliensis): Fomes root rot, where root rot incidence is low, retention of all timber and stumps from old stand being prepared for replanting will improve soil conditions, 295

Rust: of bean, Uromyces phaseoli was discovered in Wisconsin bean fields for first time in 1973, 330

---: cedar-apple rust: good control achieved in greenhouse tests on apple with triforine, 587; and quince rust, field susceptibility of 41 apple cultivars, 696

---: comandra blister rust, Pinus nigra seedlings are susceptible, 616

---: crown: the early teliospore formation by Puccinia coronata avenae on oat line 66B1285, 467

---: of flax, caused by Melampsora lini, new race was discovered, 311

---: fusiform: Cronartium fusiforme was pathogenic to sawtooth oak, an introduced species from the Orient, 198; in Western United States three native pine species, Pinus jeffreyi, P. radiata, and P. ponderosa and California black oak (Quercus kelloggii) are highly susceptible, 497; C. fusiforme applied to slash pine by telia-bearing oak leaves or by basidiospore inoculation gave similar results, 584; incidence in the South in slash and loblolly pine plantations has increased and has a wider intensified distribution since the last Southwide evaluation made 35 years ago, 1137

---: leaf: of wheat, single application of BT fungicide can increase forage yields of winter wheat by leaf rust control, 272; leaf and stem rust of wheat, development of Puccinia graminis and P. recondita near a wind barrier showed significant differences, 459; leaf and stem, of wheat, existence of different leaf rust virulences in the southern and northern populations in the United States, 806

---: of poplar: caused by two rust species, Melampsora medusae and M. larici-populina, has recently appeared in New Zealand, 423; reached epidemic proportions in experimental plots in Wisconsin, 721

---: Puccinia antirrhini, association of bacterial colonies with uredospores of P. antirrhini, 733

---: Puccinia polysora, was widespread on corn in North Carolina in 1972 and 1973, 532

---: Puccinia pygmaea, is pathogenic to red fescue, 1028

---: stem: oat stem rust, 996 isolates were identified at Cereal Rust Laboratory, 605; of wheat, prevalence of races of Puccinia graminis f. sp. tritici in the United States in 1973, 608; the detached-leaf method of culturing rust fungi may be useful for preliminary selection of wheat varieties that rust slowly in the field, 1085

Rusty spot: of peach, was controlled in New Jersey with HOE 2873, 254

Safflower (Carthamus tinctorius): assay method is described for identification of Verticillium-diseased seed lots, 926

Sand hill decline (see also Young tree decline): of citrus, 82 affected trees were indexed for virus content, 35; of unknown cause, possible diagnosis by measurement of water uptake, 801

Saskatchewan: 342

Scab: of apple: a temperature-leaf wetness recorder was built to determine infection periods for apple scab under orchard conditions, 133; caused by Venturia inaequalis, field susceptibility of 51 apple cultivars for apple scab and powdery mildew, 625; tolerance of the apple scab fungus to benzimidazole fungicides in Australia, 886

---: of peach, benomyl and BAS 3201 F

were better than other fungicides tested for control, 208
---: of pecan, benomyl and thiophanate-methyl provided control, 687
Scale: beech scale, its distribution in Pennsylvania in relation to Nectria coccinea var. faginata and Xylococculus betulae, 718
Scaphoideus luteolus: 387
Schizaphis graminum: 472
Scleroderris lagerbergii (Gremmeniella abietina): 892; 1097
Sclerophoma pythiophila: 653
Sclerophthora macrospora: 848
Sclerospora sorghi: 90; 285; 430; 825
Sclerotinia homeocarpa: 514
--- sclerotiorum (= Whetzelinia sclerotiorum): 59; 693; 700; 782
Sclerotium cepivorum: 6
Scolytus multistriatus: 382
Seed: inoculation of barley for Helminthosporium stripe infection, 32; delayed harvest of soybean seed in Indiana resulted in deterioration of soybean seed quality, 130; presence of internally-borne Diaporthe phaseolorum var. sojae in soybean seed from different locations in the U.S., 173; long smut detected on imported sorghum seed collected in Africa and India for germplasm and breeding purposes, 810
---: treatment: captan-treated soybean seeds from Illinois showed improved germination over untreated seed, 50; TCMTB-60W, as a seed treatment plus a soil treatment, controlled seedling and pod rot disease of peanut in Egypt, 176; thiabendazole-treated winter wheat seed gave fair to good control of dwarf bunt in Utah, 743; BAY 22555, as a seed dressing, gave outstanding control of seedborne bunt of wheat, 845; a seed slurry treatment of benomyl and captafol for 'Florida Sweet' corn significantly improved plant stand, 922
Seedborne infection: seedborne bean common mosaic virus causes losses in Morocco, 307; effect of cultivar and growing region on internally seedborne fungi in soybean, 332; seedborne pathogens in soybean seed lots from three locations in Illinois, 411; of Diplodia zeae and β-Phomopsis on corn seed, 414; occurrence of Phomopsis on corn, 416; Cercospora oryzae and Trichoconis padwickii in rice, Drechslera sorokiniana in pepper and flax, and Fusarium moniliforme in oats, pepper, soybean, flax and bean are new records for seedborne organisms from Argentina, 507; Exserohilum prolatum was found in corn seed imported from Guatemala and this indicates a hazard in the movement of seed from country to country when seed treatments commonly used for corn do not kill fungi within the seed, 612; seedborne Diaporthe phaseolorum var. sojae was significantly reduced in seeds from plants sprayed with benomyl, 760; D. phaseolorum and Phomopsis were isolated from immature soybean plants, 978
Seedling disease: benomyl treatment of red oak seedlings gave protection against the oak wilt pathogen, Ceratocystis fagacearum, 65; damping-off and seedling wilt of watermelon controlled with fungicides, 114; production of sugarbeet seedlings for sequential studies of Aphanomyces cochlioides disease development, 157; root rot of black walnut in Kentucky, caused by Cylindrocladium scoparium, 188; corn leaf blight pathogens Helminthosporium turcicum, and H. maydis races O and T reduce leaf and root weights in the seedling stage and differences attributable to inbred lines or to pathogens can be measured, 219; H. turcicum, and H. maydis races T and O caused reduction in top and root weight of corn seedlings, 313; Fusarium solani and Volutella sp. induce canker formation on Maesopsis eminii tree seedlings in Uganda, 463; absorption of streptomycin sulfate in soybean seedlings caused phytotoxicity, 534; injury to Pinus banksiana seedlings from Discosia pini, 677; infection of yellow poplar seedlings by Fusarium solani, 754; Mexican lime, tristeza virus symptoms were suppressed by warm temperatures, 757
Septoria apiicola: 232
--- nodorum: 21
Sequoia gigantea (giant sequoia): Fomes annosus is associated with loss of giant sequoias, 478
Short life: of peach trees, disease appears to be closely associated with site, nematodes, bacterial canker, and cold injury, 76
Silica gel: suggested as the most useful method for long-term storage of Helminthosporium maydis stock cultures, 334
Silver leaf disease: of peach, caused by Stereum purpureum, Trichoderma viride reduced symptoms, 147
Smut (Ustilago scitaminea): of sugarcane, unusual symptoms were observed on sugarcane in Hawaii, 401; Ustilago striiformis controlled on Kentucky

bluegrass with a single fall application of benomyl and PCNB, 573; long smut, Tolyposporium ehrenbergii, was detected on imported sorghum seed, 810
Snapdragon (see Antirrhinum)
Snowmold: late fall mowing to remove excessive vegetative growth of winter wheat predisposed plants to severe snowmold in Indiana, 432
Sodium azide (see Chemicals)
Soil: amendments, benomyl or thiophanate, when applied as a soil amendment, reduced ozone injury on Poa annua, 41; virgin soil or pathogen-free soil for control of bacterial wilt of greenhouse-grown tomatoes, 86; pH had little influence on resistance and susceptibility of tomato and alfalfa to Meloidogyne hapla, 594; Rhizoctonia solani infection on seed potatoes with low numbers of sclerotia decreased with increased temperatures; with high levels of inoculum there was little difference, 646; the role of temperature on effectiveness of contact nematicides for control of Pratylenchus penetrans, 708; moisture, and its effect on survival of Macrophomina phaseolina in soybean stems in soil, 1034; soil properties and fungal populations in a young-growth western hemlock stand infected with Fomes annosus, 1077
---: assay: for soilborne inoculum potential for gladiolus Fusarium disease, 184; different geographic areas of North Carolina were assayed for distribution of plant-parasitic nematodes and also for the influence of previous crops on populations, 991
---: treatments: for fungi: white rot of onion, caused by Sclerotium cepivorum, was controlled by several fungicides when dusted in furrow, 6; benomyl and thiophanate were effective in reducing injury from ozone on annual bluegrass, 41; incorporation of benomyl and chloroneb into nonvirgin soil in Florida controlled damping-off and seedling wilt of watermelon, 114; injection of benomyl into soil surrounding large nursery elms greatly reduced Dutch elm disease symptom development in inoculated twigs and mostly prevented tree mortality, 261; ETMT and BAY 22555 controlled Phytophthora cinnamomi that causes root root of Chamaecyparis and Eucalyptus, 437; benomyl injection into soil 1 month before American elms were inoculated with Ceratocystis ulmi inhibited the colonization by the fungus, 511; Fusarium wilt in Kentucky bluegrass increased with high spring fertilization, 514; benomyl + PCNB gave excellent control of stripe smut and melting out disease of Kentucky bluegrass with single fall application, 573; drench applications of ETMT controlled a new crown rot and wilt of baby's breath under greenhouse conditions, 669; fumigation with methyl bromide of a wheat field infested with soilborne wheat mosaic virus did not eliminate the vector, Polymyxa graminis, from soil, 878
---: ---: for nematodes: increased yield and marketable grade of sweetpotatoes resulted from nematicide treatments of soil infested with Meloidogyne incognita, 239; effect of crop sequence and fallow on populations of Rotylenchulus reniformis in fumigated and untreated soil in a sweetpotato nematode control program, 259; control of Rotylenchulus reniformis on cotton with nematicides, 337; fumigation of Florida potato fields with early dying disease with DD reduced nematode populations and reduced levels of Verticillium albo-atrum in plant samples, 374; 1,3-D placed deep at high dosage gave good control of nematodes in replant vineyard soil, 562; fumigants DD and DBCP in general gave better control than contact nematicides for control of Hoplolaimus columbus on soybeans in South Carolina, 641; aldicarb and oxamyl controlled root-lesion nematodes on greenhouse roses, 730; DD, EDB, ethoprop, carbofuran, fensulfothion, and phenamiphos controlled root-knot nematodes and increased yield and grade of cantaloup, 746; nematicide treatments controlled Meloidogyne incognita and increased yields of lima bean, 749; nematicides controlled Meloidogyne hapla in peony, 860; DBCP drenches were effective in killing Longidorus macrosoma in rose beds, 913; applications of UHF electromagnetic energy in combination with soil fumigation controlled the reniform nematode and increased yield in cowpea, 985; DD fumigation controlled nematodes and increased sweetpotato yields in Trinidad, 1048
Soil rot: of cucumber, caused by Rhizoctonia solani, controlled with sodium azide, 636
Sorghum: first documented observation of zonate leaf spot, caused by Gloeocercospora sorghi, on grain sorghum in Nebraska and Kansas, 267; a sap-transmissible isolate from sorghum

and/or johnsongrass in Hawaii was identified as maize dwarf mosaic virus-A, 495; quarantine inspectors found Tolyposporium ehrenbergii (long smut) on Sorghum bicolor seeds and heads collected in Africa and India for breeding purposes; it is not a pathogen in this country, 810; inoculation technique for assessing resistance in corn and sorghum varieties to Sclerospora sorghi, 825; red stripe disease of sorghum in Australia is caused by the johnsongrass-infecting strain of sugarcane mosaic virus (SCMV-J) and is related serologically to maize mosaic, 832
South Australia: 886
South Carolina: 67; 844; 1000
South Dakota: 1061
Southern corn leaf blight (see Blight)
Southern pea (see Cowpea)
Soybean: potted plants were protected against ozone when grown in soil containing 9.5 ppm of carboxin, 162; a survey was made of the most common pathogenic fungi, nematodes and endomycorrhizal fungi associated with soybean roots in Florida, 169; reactions of 70 lines and varieties of Glycine spp. to bean pod mottle virus, 191; streptomycin sulfate to control bacterial diseases was absorbed by roots and translocated in seedlings but was phytotoxic, 534; seven of 17 varieties tested showed resistance to Whetzelinia sclerotiorum, 693; reaction of cultivars to powdery mildew, caused by Microsphaera sp., 723; effect of soil moisture on survival of Macrophomina phaseolina in stems in soil, 1034
---: nematodes: low yields of cotton and soybean in east Georgia, attributed to lance nematode, may be due to other factors, 369; control of the Columbia (lance) nematode on soybeans with nematicides, 641
---: Phytophthora megasperma var. sojae: a more virulent strain of P. megasperma var. sojae than race 3 has been recovered from diseased soybeans, race 4 is proposed, 352; 95 of 266 soybean germplasm strains were resistant to all four races of P. megasperma var. sojae, 789; effect of Rhizobium japonicum and Endogone mosseae and soybean root rot, caused by Pythium ultimum and P. megasperma var. sojae, 221
---: seedborne infection: Diaporthe phaseolorum var. sojae was the predominant organism isolated from soybean seed in Illinois, 50; seed infection with Diaporthe phaseolorum var. sojae and Alternaria spp. increased with delay in harvest, 130; occurrence of Diaporthe phaseolorum var. sojae in seed lots was variable with harvest year, location and variety in seeds from different states in the U.S., 173; growing region and cultivar affect the percentage of internally seedborne fungi and Aspergillus melleus pathogenicity in soybean, 332; seed lots of 'Amsoy' and 'Wayne' soybean from three different regions of Illinois were tested for internally-borne fungi, 411; Fusarium moniliforme detected on soybean seed in Argentina, a new record, 507; benomyl sprays reduced internally seedborne fungi in soybean seeds, 760; isolation of seedborne Diaporthe phaseolorum and Phomopsis, 978
Spain: 147; 442
Sphaeronema sp.: 180
Sphaerotheca fuliginea race 2: 899
Spiroplasma (see also Mycoplasma): spiroplasma filaments were detected in corn stunt-infected individuals of the vector leafhopper Dalbulus maidis, 1109
--- citri: 797
Spore collection: a vacuum-filter method for collecting ascospores of Whetzelinia sclerotiorum, 190
Spore production: induction of early teliospore formation in oat rusts, 467
Sporulation: of Helminthosporium maydis race T in relation to relative humidity, 297; of Glomerella cingulata, cause of dieback of camellias, 300
Squash (see Cucurbits)
Sri Lanka: 293
Stem pitting: of peach, found in 2-year-old commercial orchard in Georgia in 1973, 553
Stereum purpureum: 71; 147
Stone fruits: method described for producing apothecia of Monilinia fructicola from brown rot-infected fruit in South Carolina where the sexual stage is rarely seen, 844
Storage: of stock cultures of Helminthosporium maydis in several ways, silica gel appears to be the most useful method of storage, 334
Strains: peanut mottle virus strain M2 and the necrosis strain were found in Georgia peanut fields, 107; five strains of tobacco etch virus and nine strains of potato virus Y were identified in commercial peppers in California, 1002; strain D of sugarcane mosaic virus occurs in Morocco, 1121
Strawberry: six new strawberry indicator clones were evaluated for detection and diagnosis of 12 graft-transmissible

viruses, 28; a modified leaf grafting technique for detection of graft-transmissible diseases of strawberry, 203
Sugarbeet: root-knot nematodes, overwintered galls contain nematodes that can reestablish infections in susceptible plants, 88; two seedling production systems were developed for studying sugarbeet-Aphanomyces cochlioides interactions, 157; benomyl-resistant strains of Cercospora beticola have appeared in Texas, 434; powdery mildew, caused by Erysiphe polygoni, reached epiphytotic proportions in California sugarbeet fields in 1974, 904; six fungicides were effective in control of Cercospora leaf spot of sugarbeet and increased root yields and sucrose content, 952
Sugarcane: nematodes of sugarcane new to Trinidad, 122; convoluted sori from lateral buds and growth anomalies associated with smut infection were found recently on sugarcane in Hawaii, 401; Xanthomonas albilineans was isolated from diseased tissue by leaf-sectioning technique, 439; sugarcane mosaic virus strain D was identified in variety N. Co. 310 in Morocco, 1121
Sunflower: diseases of sunflower in Tanzania, 896
Survey: of the most common pathogenic fungi, nematodes and endomycorrhizal fungi associated with soybean roots in Florida, 169; of corn stalk rot in Minnesota, 363; of cotton and soybean fields in east Georgia for lance nematodes, 369; of seedborne organisms of Argentina, 507; of beech bark disease distribution in Pennsylvania in 1973, 718
Susceptibility: of gladiolus cultivars to soilborne Fusarium in field samples, 184; of Phaseolus lathyroides to soybean mosaic virus, 322; susceptibility ratings of 51 apple cultivars for apple scab and powdery mildew were calculated from a survey of 37 research and extension workers throughout the United States and Canada, 625; of 41 apple cultivars to cedar-apple rust and quince rust, 696; of 46 apple cultivars to fire blight, with ratings for three geographic climatic regions in North America, 819; of cucumber accessions to viral diseases, 735; differences in nematode damage among clonal apple rootstocks, 1007; of red raspberry cultivars to spur blight, 1024
Sweetpotato: cellular inclusions and viruslike particles found in sweetpotato roots having the "hardcore" condition, 17; soil treatments with DD, EDB and BAY 68138 (phenamiphos) increased sweetpotato yield and marketable grade, and controlled Meloidogyne incognita; carbofuran gave best control of sweetpotato flea beetle and southern potato wireworm, 239; Rotylenchulus reniformis decreased in fumigated plots or untreated plots planted to corn, but increased in fumigated and untreated plots planted to tomato, bodie beans or sweetpotato, 259; nematode control with DD fumigation increased sweetpotato yields in Trinidad, 1048
Syagrus calcaratus: 616
Sydowia polyspora: 94
Symbiosis: of Fusarium fungi and ambrosia beetles, is thought to be the cause of girdling canker of black walnut, 1044
Symptomatology: of seedborne bean common mosaic virus, 307
Synchytrium endobioticum: 291

Tanzania: 896
Tea: leaf blister blight, caused by Exobasidium vexans, superiority of copper fungicides over organic fungicides appears to be due to the excellent tenacity of the copper under monsoon conditions, 928
Temperature: air, recorded simultaneously with wetness duration on the same drum for immediate evaluation of data for apple scab infection periods, 133; higher soil temperatures decreased Rhizoctonia solani infection on seed potatoes with low numbers of sclerotia, 646; 82-104°F and 78-80° daytime and nighttime temperatures totally suppressed tristeza symptoms in Mexican lime for six isolates, 757; tree temperatures of healthy and young tree-declined Valencia sweet orange budded on rough lemon rootstock measured by remote sensing were not feasible for previsual diagnosis of diseased trees, 793; controlled temperature chamber with humidity maintained at 100% continuously can be programmed for producing reproducible infections caused by many different pathogens, 814; warm glasshouse temperatures 28 to 40°C daytime max and 25.6°C nighttime minimum for 3 to 4 months inactivated several citrus viruses, 850
Tennessee: 1073
Texas: 23; 80; 504; 773; 780; 882
Thielaviopsis basicola: 54; 490
Thistle (see Canada thistle)
Tilletia caries: 845
--- controversa: 743
--- foetida: 845
Tip dieback: Sydowia polyspora found on

white fir twigs for first time in California, 94
Tobacco: black shank of tobacco found in Connecticut in 1973, 428
---: air pollution: plants growing in soil containing 9.5 ppm carboxin were protected from ozone injury, 162; leaf injury from air pollutants, 683
---: virus: tobacco vein mottling virus was found to be distributed throughout Madison County, N.C., the leading producer of burley tobacco in N.C., 521; tobacco etch virus, 'V 20' tobacco was found to be immune to six of seven isolates of TEV; it may be useful in breeding for resistance, 658; tobacco etch virus, strain differentiation of isolates based on resistance and susceptibility of pepper cultivars, 1002; tobacco vein mottling virus is found in Tennessee, 1073
Tolerance: walnut hybrids (Juglans hindsii x J. regia) were more tolerant of root-lesion nematodes than other Juglans species, 630
Tolyposporium ehrenbergii: 810
Tomato: fern leaf, yields of plants growing adjacent to fern leaf-dwarfed plants were no greater than yields of healthy plants two rows away, 161; big bud, found in New York State, 211; vertical distribution of Fusarium oxysporum f. sp. lycopersici race 2 to a depth of 90 cm in greenhouse soil may require modification of growing practices in greenhouses until the release of a resistant cultivar, 320; Lycopersicon: Fusarium host: pathogen disease resistance development in relation to horizontal and vertical resistance, 579; Rhizoctonia soil rot resistance, segregation of F_2 population of the cross (Merit (susceptible) x PI 193407 (resistant)) was scattered over entire range of disease shown by parents, suggesting resistance is polygenic, 406; brown root rot, caused by Pyrenochaeta lycopersici, is reported for first time in North America on tomato, 483; resistance and susceptibility of tomato varieties to Meloidogyne hapla was not particularly influenced by soil pH, 594
---: air pollution: protection from ozone injury with carboxin, 162; PAN-type injury symptoms are reported for the first time on vegetation in Canada, 1105
---: bacterial wilt: a virgin-soil technique for controlling wilt in greenhouse-grown tomatoes, 86; Pseudomonas solanacearum confirmed in Guam as organism responsible for bacterial wilt infection, 793
---: virus: tobacco mosaic virus, causes severe losses in tomato, the most important vegetable crop in Morocco, 309; infection of virus elimination plots and home gardens with an apparently new destructive viruslike disease in Washington, 965
Toxin: several oat cultivars were tested for a relationship between calcium nutrition and sensitivity to a wilt toxin, victorin, produced by Helminthosporium victoriae, 3
Transmission: of a mosaic disease (identified as tobacco mosaic virus) of white ash to woody and herbaceous hosts, 536; of strains of grapevine fanleaf virus by Xiphinema index, 549; Artichoke Italian latent virus is apparently not seed-transmitted, 691
Tree breeding: for resistance to poplar leaf rust, 721; for resistance to Dutch elm disease, 727; 784
Trichoconis padwickii: 507
Trichoderma viride: in large quantities, ungerminated spores of T. viride inhibited growth of Stereum purpureum propagules in plate culture, 71; was effective on plum trees in preventive control of silver leaf disease, caused by Stereum purpureum, 147; heat-killed spores inhibited in vitro mycelial growth of Botrytis cinerea, Monilinia laxa, M. fructigena, and Phomopsis viticola, 915
Trichodorus sp.: 991
--- spp.: 169
--- allius: 269
--- christiei: 374; 822
--- minor: 122
--- teres: 269
Trinidad: 122; 1048
Tsuga heterophylla (western hemlock): reported occurrence of Rhizina root rot in an old-growth conifer stand in the Pacific Northwest, 492; preliminary data suggest that soil in vicinity of Fomes annosus-affected trees is lower in organic matter and has a lower pH, than soil around healthy trees, 1077
Tubercularia ulmea: 937
Tulip: mycoplasmalike bodies were found in phloem sieve elements in malformed tulips in Israel, 1142
Tylenchorhynchus sp.: 991
--- spp.: 1048
--- claytoni: 374; 822
--- martini: 122
Tylenchulus semipenetrans: 882
Typhula itoana: 432

Ulmus americana (American elm, see Elm)
Ultra-high frequency (UHF) electromagnetic energy: UHF in combination with 1,3-D controlled reniform nematode

and common purslane in field of southern peas, 985
Uncinula necator: 486
Urd (see Vigna radiata)
Uromyces phaseoli: 330
Ustilago scitaminea: 401
--- striiformis: 573

Variety development: concepts for the overall theory of disease control are outlined and discussed with respect to the imposition of federal controls on breeding programs, 579
Vectors associated with: new virus of corn from Venezuela, Peregrinus maidis, 122; potato virus Y in peppers, Myzus persicae, 141; tobacco rattle virus, Trichodorus allius, and T. teres nematodes, 269; Dutch elm disease, Scolytus multistriatus, 382; elm phloem necrosis, Scaphoideus luteolus, 387; barley yellow dwarf virus, Rhopalosiphum padi, R. maidis, Macrosiphum avenae, and Schizaphis graminum, 472; maize dwarf mosaic virus strain A, Rhopalosiphum maidis, 495; grapevine fanleaf virus, Xiphinema index, 549; newly described mosaic disease of okra, is readily transmitted by two chrysomelid beetles, Syagrus calcaratus and Podagrica uniformis, 616; carrot motley dwarf virus, Cavariella aegopodii, 766; rice tungro virus, Nephotettix virescens, 856; soilborne wheat mosaic virus, Polymyxa graminis, 878; new viral disease of wheat in India, Cicadulina mbila, 968; an aphid-borne virus disease of irrigated cowpea in Nigeria, Aphis craccivora, 1080; sugarcane mosaic virus, Rhopalosiphum maidis, 1121; artichoke latent virus, Myzus persicae, 1123
Venezuela: 122; 1032
Venturia inaequalis: 133; 625; 886
Verticillium sp.: 180
--- albo-atrum: 373; 374
--- dahliae: 926; 996
--- malthousei: 987
Victorin: oat cultivars tested to determine whether a relationship exists between calcium nutrition and sensitivity to this wilt toxin, 3
Vigna mungo (urd): a susceptible host of Heterodera vigni, 345
--- radiata (mung): host of Heterodera vigni, in India, 345
--- unguiculata ssp. unguiculata (see Cowpea)
Viola odorata (sweet violet): is natural host of bean yellow mosaic virus, 155
--- tricolor: new host of tobacco streak virus, 354
Virginia: 150; 396; 455; 456
Virus: of sweetpotato, viruslike particles and cellular inclusions found associated with "hardcore" of sweetpotato roots, 17; of Eucalyptus, viruslike particles were isolated from diseased trees in Australia, 92; of corn, a new virus found in Venezuela causes stunting and white to yellow stripes on leaves, 122; a technique for purifying viruses in laboratories where ultracentrifuges are not available, 126; a previously undescribed mosaic disease of okra found in Nigeria, 616; of Hibiscus rosa-sinensis, a new ringspot virus found in Nigeria, 1040; aphid-borne virus disease of cowpea in Nigeria is either strain of cowpea mosaic virus or a new virus not previously described, 1080
---: of artichoke: purification, electron microscopy and serology of artichoke Italian latent virus, 691; artichoke mottled crinkle virus, found in artichoke in Morocco, 1117; artichoke latent virus is widespread in commercially grown artichokes in Morocco, 1123
---: barley yellow dwarf virus, mixed infections of BYDV isolates were recovered from winter wheat and barley in New York, 472
---: bean common mosaic virus, is one of principal causes of yield loss in green snap bean production in Morocco, 307
---: bean pod mottle virus, 70 lines and varieties of Glycine spp. were tested for resistance to this virus, 191
---: bean yellow mosaic virus, Viola odorata is a natural host, 155
---: Bidens mottle virus in lettuce and escarole fields in Florida, 1087
---: carrot motley dwarf, found to occur in both Oregon and Washington, 766
---: of citrus: primarily three transmissible agents: exocortis, tristeza and a stem pitting factor were found in indexed trees having young tree decline or sand hill decline, 35; exocortis, of citron, thin-layer chromatography may be helpful in early diagnosis of exocortis disease, 82; Satsuma dwarf virus, Physalis floridana is a good production host, 603; tristeza, very warm temperatures suppressed symptoms of tristeza in Mexican lime, 757; psorosis, concave gum, infectious variegation, and vein enation and tristeza viruses were inactivated by holding at warm glasshouse temperatures for 3 to 4 months, 850
---: cucumber mosaic virus, that causes fern leaf of tomato, its effect on yield, 161

Virus: dasheen mosaic virus, was identified in Venezuela in dasheen (Colocasia esculenta) and yautia (Xanthosoma spp.), 1032
---: grapevine fanleaf virus, three strains of the virus are apparently equally available to the vector, Xiphinema index, and acquisition access threshold time is less than 5 minutes, 549
---: lettuce mosaic virus (LMV): indexing lettuce seeds for LMV, airstream separation of light and heavy seeds concentrated the virus-containing seeds in the light portion, 1037; lettuce mosaic and Bidens mottle viruses in lettuce and escarole fields in Florida, 1087
---: maize chlorotic dwarf virus, was identified from diseased johnsongrass in southern Illinois, 420
---: maize dwarf mosaic virus, (MDMV), a sap-transmissible virus from sorghum and/or johnsongrass from the Hawaiian Islands was identified as MDMV strain-A, 495
---: maize mosaic, the viruses responsible for red stripe disease of sorghum and mosaic of maize were similar and were found to be serologically related, and were caused by the johnsongrass-infecting strain of sugarcane mosaic virus (SCVM-J), in Australia, 832
---: onion yellow dwarf is one of the most serious diseases of onions in Morocco, 252
---: peanut mottle virus, a survey of 117 peanut fields in Georgia showed an average incidence of 26% in 1973, 107
---: peanut stunt, on white clover, first report of this virus in Florida, 830
---: of pepper: potato virus Y is the most serious disease of pepper in Morocco, 141; potato virus Y and tobacco etch viruses were isolated most frequently from peppers in California, 1002
---: potato virus Y: 141; 1002
---: red stripe disease of sorghum, is caused by the johnsongrass-infecting strain of sugarcane mosaic virus (SCMV-J) and is serologically related to a virus isolate causing maize mosaic in Australia, 832
---: rice tungro virus, Paspalum distichum, Echinochloa colonum, Cyperus rotundus, and Cynodon dactylon serve as alternate hosts of virus, 856
---: soybean mosaic virus, a potential forage plant (Phaseolus lathyroides) in Brazil is susceptible to the virus, 322
---: squash mosaic virus, resistance in two accessions of "jelly melon," 735
---: of strawberry: six new indicator clones were tested for use in detection and diagnosis of 12 graft-transmissible viruses, 28; graft-transmissible diseases, of strawberry, 203
---: sugarcane mosaic virus: johnsongrass-infecting strain is cause of red stripe disease of sorghum in Australia, 832; strain D was identified in Morocco, 1121
---: sweet cherry raspleaf, a survey shows CRLV is becoming an important disease in Colorado sweet cherry orchards, 26
---: tobacco etch virus, resistance in 'V 20' tobacco to six of seven isolates, 658; in peppers in California, 1002
---: tobacco mosaic virus, was identified as the cause of a mosaic disease of white ash; it is the first evidence that TMV causes a disease of a tree, 536
---: tobacco rattle virus, Trichodorus allius and T. teres are vectors of three tobacco rattle virus isolates affecting potato production in Oregon, 269
---: tobacco streak virus, five new experimental hosts have been found, 354
---: tobacco vein mottling virus: distribution and incidence in Madison County, North Carolina, 521; found in Tennessee affecting burley tobacco, 1073
---: of tomato: tomato spotted wilt virus, this is the first report of this virus on peanut in the United States, in Texas, 23; fern leaf of tomato, caused by cucumber mosaic virus; healthy plants adjacent to dwarfed diseased plants showed no yield increase over healthy plants two rows away, 161; tobacco mosaic virus caused severe losses in tomatoes in Morocco, 309; a viruslike disease of tomato was found in curly top virus elimination plots in Washington, 965
---: watermelon mosaic virus (WMV): WMV-2, serious losses in cucurbits in Morocco, 143; three watermelon varieties were evaluated for effect of WMV on growth habit and fruit production, 195; WMV-2 found in Chile on squash, honeydew melon, and zucchini squash, 599; WMV-1 resistance in Cucumis metuliferus, 735
---: of wheat: wheat spindle streak mosaic, 10 wheat cultivars were evaluated for its incidence and its effect on yield, 522; soilborne wheat mosaic virus (SBWMV) has been spread in Florida, possibly because its vector Polymyxa graminis has been spread by farm machinery, 763; resistance to SBWMV found in wheat cultivars, but in Florida cultivars need resistance to other prominent diseases as well, 878; a

new viral disease with striate symptoms was found on several wheat and barley cultivars in India, 968
Volutella sp.: 463

Walnut (Juglans): first reported occurrence of Cylindrocladium scoparium root rot in Kentucky, on black walnut seedlings, 188; all Juglans species tested were very susceptible to root-lesion nematode disease, 630; a stem girdling canker of black walnut found in Midwest is thought to be caused by a Fusarium-ambrosia beetle complex, 1044
Washington: 6; 10; 62; 554; 701; 766; 965
Water hyacinth (Eichhornia crassipes): leaf spot, caused by Cercospora piaropi, was recorded for the first time in the United States in Florida, 277
Watermelon (see Cucurbits)
Weather: uncommonly cool spring is thought to be responsible for severe outbreak of oat leaf blotch in Texas, 80
West Virginia: 291
Wheat: glume blotch, caused by Septoria nodorum, in Georgia was more severe on early cultivars, 21; snowmold, late fall mowing predisposed plants to severe snowmold, 432; Ascochyta sorghi was collected from wheat cultivars in Virginia, 455; earlier seeding of winter wheat, protection with benomyl spray, and wheat resistant varieties should increase erosion control and limit Cercosporella foot rot in Washington, 554; Cephalosporium stripe of winter wheat and barley in Kansas, 566; BAY 22555 gave excellent control of seedborne bunt in Australia, but was ineffective on soilborne bunt, 845; Pseudomonas syringae incited a leaf necrosis on spring and winter wheats in South Dakota, 1061
---: rust: leaf rust, caused by Puccinia recondita f. sp. tritici, 4-n-butyl-1,2,4-triazole (BT) is an ideal fungicide for measuring the effects of leaf rust on yields of winter wheat forage, 272; leaf and stem rust, near a wind barrier, differ in their development, 459; leaf and stem rust, control of rusts with zinc + maneb spray increased yields in test plots up to 44%, 469; stem rust, caused by Puccinia graminis f. sp. tritici, the prevalence of races present in the U.S. in 1973, 608; leaf and stem rust, major differences were detected between the southern (Beeville, Texas), and northern (Rosemount, Minnesota) rust populations, 806; slow rusting of wheat varieties in the field is correlated with stem rust severity on detached leaves in the greenhouse, 1085
---: virus: barley yellow dwarf virus, mixed infections of the virus were common in winter wheat in New York, 472; effect of wheat spindle streak mosaic on 10 winter wheat cultivars, 522; soilborne wheat mosaic virus, its presence, distribution and yield loss in Florida, 763; cultivars with resistance to soilborne wheat mosaic virus are available in Florida but multiple disease resistance is necessary, 878; viral-like particles were found in phloem of wheat and barley with striate symptoms, 968
Whetzelinia sclerotiorum (Sclerotinia sclerotiorum): 190; 379; 693; 700; 782
White mold: Whetzelinia sclerotiorum infection in 'Great Northern' dry beans was reduced in trellis-grown plants in comparison with pruned "runners," or 2,3,5-triiodobenzoic acid (TIBA)-sprayed plants, 379; of soybean, caused by W. sclerotiorum, several varieties show resistance, 693; tolerance to white mold infection was found in several dry edible bean varieties grown in Nebraska, 782
Wilt: bacterial: of tomatoes, control with soil that doesn't contain the pathogen, Pseudomonas solanacearum, 86; of tomato and eggplant, is distributed throughout Guam, 793
---: crown rot and wilt of baby's breath (Gypsophila paniculata), caused by Phytophthora parasitica, in Florida, 669
---: Fusarium: of muskmelon, was recorded for first time in Washington, caused by Fusarium oxysporum f. sp. melonis, 10; common wilt of pea, caused by Fusarium oxysporum f. sp. pisi, was found in Washington in field planted to a susceptible pea variety, 62; seedling wilt of watermelon, caused by Fusarium and Rhizoctonia, controlled with fungicides, 114; of tomato, caused by F. oxysporum f. sp. lycopersici, race 2-infested soil in Ohio is a threat to greenhouse growers, 320; of tomato, as a disease model for evaluation of concepts of variety development and disease control with host resistance, 579
---: Goss's or Nebraska wilt (see Leaf freckles)
---: oak: caused by Ceratocystis fagacearum, colonization was prevented in red oak seedlings by benomyl drench, 65; oak wilt fungus identified in two counties in Missouri, 92; restricted spread of oak wilt in Arkansas is possibly caused by a natural biological control, a competing fungus, Hypoxylon atropunctatum, 622;

in red oak trees, disease incidence was greater and the incubation period was more uniform in trees bole-inoculated than root-inoculated with Ceratocystis fagacearum, 738
---: vascular wilt of aster, caused by Phialophora fastigiata, 57
---: Verticillium: Verticillium wilt infection in safflower seeds was detected by seed assay, 926; wilt of cotton, reported for the first time from Iraq, 996
Wind barrier: in a wheat leaf and stem rust development study, a wind barrier had significant influence on incidence and severity of pathogens, 459
Windthrow: of aspen, loss by windthrow in a Colorado aspen stand was a result of the predisposition action of Fomes annosus, 110
Wisconsin: 226; 229; 330; 653; 941
Witches' broom: of pigeonpeas in Puerto Rico, 96
Wood decay fungi: benomyl-malt agar used for the purification of cultures of wood decay fungi in the process of transferring and maintaining cultures, 902

Xanthomonas albilineans: 439
--- begoniae: 814
--- malvacearum: 373
--- phaseoli: 278
Xanthosoma spp. (yautia): dasheen virus was found in plants of Xanthosoma in Venezuela, 1032
Xiphinema sp.: 991
--- spp.: 169
--- americanum: 633
--- index: 549; 562
Xylococculus betulae: 718

Yautia: infection with mosaic virus in Venezuela, 1032
Yellow poplar (Liriodendron tulipifera): infection with Cylindrocladium floridanum in New Zealand, 705; Fusarium solani infected yellow poplar seedlings through wounds on stems and through roots whether wounded or not, 754
Yellow tuft: of bluegrass and bentgrass, Sclerophthora macrospora reported for the first time associated with this disease, 848
Yield: effect of fern leaf on the yield of tomatoes, the farmer would gain 0.64% yield for every day his plants escape infection with fern leaf, 161; of 'Great Northern' beans, yield was affected by the modified architecture; trellis support resulted in higher bean yields than by pruning or by a growth regulator, and there was less white mold infection, 379
Yield increase: of winter wheat forage with single application of BT fungicide, 272; wheat yields on test plots were increased up to 44% by controlling leaf and stem rust development on selected cultivars with zinc + maneb spray, 469; cantaloup yield and grade were increased by nematicidal control of Meloidogyne incognita, 746; lima bean yields were increased by control of Meloidogyne incognita with nematicides, 749; of sweetpotato in Trinidad after fumigating nematode-infested soil, 1048
Yield loss: of cotton in 1973 from parasitic diseases, 373; of Concord grapes following sprays with cupric hydroxide, 486; seed treatment with thiabendazole at 8 oz/100 lb rate caused yield reduction of winter wheat, but controlled dwarf bunt, 743; from soilborne wheat mosaic virus in Florida, 763; of beans in Kenya from root-knot nematodes, 1020
Young tree decline (YTD): of citrus, in Florida, 35; attempt at previsual diagnosis of disease by use of a remote sensing infrared thermometer was not a reliable indicator, 793; restricted uptake of water injected into tree trunks of diseased trees may prove to be a useful diagnostic tool for identifying this kind of decline, 801; 3 of 5 YTD-affected trees given 7 treatments with tetracycline over a 20-month period showed noticeable improvement, the other two did not, 895; soil and spray treatments and erythorbic acid, sodium erythorbate, ascorbic acid and/or sodium ascorbate resulted in renewed growth and recovery from YTD symptoms in 'Marsh' grapefruit and 'Valencia' oranges, 918; see also Blight: citrus, 1051

Prepared by Marguerite G. Benedict

U.S. GOVERNMENT PRINTING OFFICE : 1975 O - 597-490

UNITED STATES DEPARTMENT OF AGRICULTURE
Agricultural Research Service
Beltsville, Maryland 20705

Official Business
Penalty for Private Use, $300

POSTAGE AND FEES PAID
United States Department of Agriculture

U.S.MAIL

PDR XEROX300XB R 1
XEROX UNIV MICROFILMS
SERIALS DEPT
300 N ZEEB RD
ANN ARBOR MI 48106

1